

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 January 2001 (11.01.2001)

PCT

(10) International Publication Number
WO 01/02430 A2

(51) International Patent Classification⁷: **C07K 14/00**

(74) Agent: WITTOP KONING, T., H.; Exter Polak & Char-
louis B.V., P.O. Box 3241, NL-2280 GE Rijswijk (NL).

(21) International Application Number: **PCT/EP00/06401**

(22) International Filing Date: **5 July 2000 (05.07.2000)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:
99202214.5 **5 July 1999 (05.07.1999) EP**

(71) Applicants (for all designated States except US):
CROPDESIGN N.V. [BE/BE]; Technologiepark 3,
B-9052 Gent (BE). **UNIVERSIDADE FEDERAL DO**
RIO DE JANEIRO [BR/BR]; CEP-21941-590 Rio de
Janeiro, RJ (BR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **HEMERLY, Adri-**
ana, Silva [BR/BR]; Rua Padre Achotegui, 60-1004,
Leblon, CEP-22430-090 Rio de Janeiro, RJ (BR). **FER-**
REIRA, Paulo, Cavalcanti, Gomes [BR/BR]; Rua Padre
Achotegui, 60/1004, Leblon, CEP-22430-090 Rio de
Janeiro, RJ (BR). **ROMBAUTS, Stephane** [BE/BE];
Nederpolder 23, B-9000 Gent (BE).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).


Published:

— Without international search report and to be republished
upon receipt of that report.

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: **PLANT PROTEINS**

CDC7 Gene Structure



(57) Abstract: The present invention relates to at least partially purified protein, capable of modulating the DNA replication in plants, muteins thereof, DNA coding therefor and to a method to confer to one or more plant cells the capacity to provide such a protein or mutein. The invention also relates to plants, comprising the said DNA and the progeny thereof.

BEST AVAILABLE COPY

Title: Plant proteins.

5

The present invention relates to at least partially purified protein, capable of modulating the DNA replication in plants, muteins thereof, DNA coding therefor and to a method to confer
10 to one or more plant cells the capacity to provide such a protein or mutein. The invention also relates to plants, comprising the said DNA and the progeny thereof.

The regulation of the cell cycle in plants is poorly understood. Most of the knowledge regarding the regulation of
15 DNA replication, also known as the S-phase of the cell cycle regulation originates from experimental data obtained in yeast and mammalian cells. However, the importance to understand the cell cycle regulation in plant cells has become increasingly important in agriculture, e.g. to control growth of plants at
20 stress conditions, to obtain resistance against parasites that block or modulate the cell cycle regulation, or to improve the yield of agriculturally important crops. Further, one might be interested to intervene in the cell cycle regulation by allowing further rounds of DNA replication, but simultaneously
25 preventing further cell cycle progress by blocking the subsequent mitosis. In this way, cells may be obtained having multiple sets of their genetic material, so that plants with a high rate of endoreduplication may be generated. The term "endoreduplication" means recurrent DNA replication without
30 consequent mitosis and cytokinesis.

From experiments in yeast, it is known that DNA replication and mitosis are coupled events in the cell cycle. Paulovich et al., 1997; Cell 88, 315-321. Genetic studies in yeast for example suggest that the CDC7 serine-threonine kinase
35 plays a role in the initiation of DNA synthesis. Evidence has been presented that CDC7 is apparently directly involved in the activation of individual early- as well as late replication origins during S-phase (Bousset and Diffley, 1998, Genes Dev 12, 480-490; Donaldson et al., 1998, Genes Dev 12, 491-501).
40 The protein levels of CDC7 are constant during the cell cycle. Activation of CDC7 as a kinase occurs at the G1/S transition of the cell cycle and is dependent on the binding with another factor, DBF4, at the G1/S transition of the cell cycle,

probably by phosphorylating proteins at the origins (Kitada et al, 1992; Genetics 131: 21-29, Lei et al; Genes and Development 11, 3365-3374, 1997). In order to function as a kinase, the CDC7 kinase may be a substrate for one or more phosphorylation events. Overexpressed kinase-negative mutants of CDC7 arrest yeast cells in the G1 to S transition and inhibit growth. Further experiments showed that the inactivation of wild-type CDC7 function probably can be explained through titration of DBF4 by the inactive cdc7 mutant proteins (Ohtoshi et al., 1997, Mol Gen Genet 254, 562-570). In addition to mechanisms to control the onset of DNA replication, other mechanisms contribute to restrict DNA replication to occur only once during the cell cycle. For example, the CDC16, CDC23 and CDC27 proteins are part of a high molecular weight complex, known as the anaphase promoting complex (APC) or cyclosome, (see Romanowski and Madine, Trends in Cell Biology 6, 184-188, 1996, and Wuarin and Nurse, Cell 85, 785-787 (1996), both incorporated herein by reference). The complex in yeast is composed of at least 8 proteins, the TPR (tetratricopeptide repeat) containing proteins CDC16, CDC23 and CDC27, and five other subunits named APC1, APC2, APC4, APC5 and APC7 (Peters et al. 1996, Science 274, 1199-1201). The APC targets its substrates for proteolytic degradation by catalyzing the ligation of ubiquitin molecules to these substrates. APC-dependent proteolysis is required for the separation of the sister chromatids at meta- to anaphase transition and for the final exit from mitosis. Among the APC-substrates are the anaphase inhibitor protein Pds1p and mitotic cyclins such as cyclin B, respectively (Ciosk et al. 1998, Cell 93, 1067-1076; Cohen-Fix et al. 1996, Genes Dev 10, 3081-3093; Sudakin et al. 1995, Mol Biol Cell 6, 185-198; Jorgensen et al. 1998, Mol Cell Biol 18, 468-476; Townsley and Ruderman 1998, Trends Cell Biol 8, 238-244). To become active as a ubiquitin-ligase, at least CDC16, CDC23 and CDC27 need to be phosphorylated in the M-phase (Ollendorff and Donoghue 1997, J Biol Chem 272, 32011-32018). Activated APC persists throughout G1 of the subsequent cell cycle to prevent premature appearance of B-type cyclins which would result in an uncontrolled entry into S-phase (Irniger and Nasmyth 1997, J Cell Sci 110, 1523-1531). It has been demonstrated in yeast that mutations in

either of at least two of the APC components, CDC16 and CDC27, can result in DNA overreplication without intervening passages through M-phases (Heichman and Roberts 1996, Cell 85, 39-48). CDC16, CDC23 and CDC27 all are tetratricopeptide repeat (TPR) containing proteins. A suggested minimal consensus sequence of the TPR motif is as follows: $X_3-W-X_2-L-G-X_2-Y-X_8-A-X_3-F-X_2-A-X_4-P-X_2$ (Lamb et al. 1994, EMBO J 13, 4321-4328; X denotes any amino acid, X_n a stretch of n of such amino acids). However, the consensus residues can exhibit significant degeneracy and little or no homology is present in non-consensus residues. The hydrophobicity and size of the consensus residues, rather than their identity, seems to be important. TPR motifs are present in a wide variety of proteins functional in yeast and higher eukaryotes in mitosis (including the APC protein components CDC16, CDC23 and CDC27), transcription, splicing, protein import and neurogenesis (Goehl and Yanagida 1991, Trends Biochem Sci 16, 173-177). The TPR forms a α -helical structure, tandem repeats organize into a superhelical structure ideally suited as interfaces for protein recognition (Groves and Barford 1999, Curr Opin Struct Biol 9, 383-389). Within the α -helix, two amphipathic domains are usually present, one at the NH_2 -terminus and the other near the $COOH$ -terminus (Sikorski et al. 1990, Cell 60, 307-317).

In order to understand the mechanisms playing a role in plant cell cycle regulation, in particular the DNA replication, and to understand endoreduplication in plants, the present inventors isolated several novel plant DNA sequences, coding for novel proteins, or novel amino acid sequences thereof involved in the modulation of DNA replication, using degenerated PCR primers based on known genomic or cDNA sequences, e.g. of yeast, mammals and insects.

"Capable of modulating the DNA replication in plants" is to be understood as the capacity of a protein to alter the natural DNA replication mechanism in the said plant, e.g. by up- or down-regulation of the DNA replication in a way, different from the natural situation, or to a higher or lower extent with respect to the natural situation. The natural situation is to be understood as the situation wherein DNA replication takes place in plants, in which the DNA replication machinery is not affected by the introduction of foreign

genetic material. Such altering includes mediating e.g. the onset of DNA replication, the rate and extent of DNA replication, the timing of DNA replication in the cell cycle, coupling or uncoupling DNA replication with/from actual
5 subsequent cell division etcetera.

Proteins

By using degenerated oligonucleotides as amplification primers, based on conserved sequence regions of the CDC7
10 homologue gene of *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe* and on conserved sequence regions of the CDC27 homologue genes of *Schizosaccharomyces pombe* and from *Aspergillus nidulans*, *drosophila* and human, the present inventors surprisingly found such novel proteins and amino acid
15 sequences. Reference is made to the examples.

Thus, novel cDNAs and proteins comprising one or more novel amino acid sequences were found. The present invention therefore relates in the first place to an at least partially purified protein, capable of modulating DNA replication in
20 plants, at least comprising in the amino acid sequence

a) one or more of the amino acid sequences chosen from the group consisting of those, given by SEQ ID NOS 2, 3 and 4,

b) one or more of the amino acid sequences chosen
25 from the group consisting of those, given by SEQ ID NOS 6, 7, 10 and 12.

c) one or more amino acid sequences having at least 50% amino acid identity with those of a), or

d) one or more amino acid sequences having at
30 least 50% amino acid identity with those of b).

By using degenerated CDC7 oligonucleotides to amplify a PCR fragment as is indicated above and will be further detailed in the examples, a novel *Arabidopsis* cDNA comprising coding sequence of an novel *Arabidopsis* CDC7 homologue gene was found
35 (SEQ ID NO 8). By comparison of the said sequences with sequences of the EMBL and EMBLnew databanks, a genomic *Arabidopsis thaliana* sequence was found (accession number Z97342). In this known genomic sequence however, only 11 exons were identified. The novel DNA according to the present
40 invention however clearly indicated the presence of 3

additional coding sequences coding for novel amino acid sequences (SEQ ID NO 2, 3, 4) being part of a DNA replication modulating plant protein, homologous to yeast CDC7.

The novel amino acid sequence SEQ ID No 2
5 (GYGIVYKATRKTDGTEFAIK) is located in two highly conserved domains in protein kinases, Domain I and II (Hawks et al., 1988, Science 241, 42-52). The sequence GYGIV is part of the nucleotide (ATP) binding domain, also known as Domain I in
10 protein kinases. Domain I is part of the catalytic domain of protein kinases. The Glycines (G) are believed to form an elbow around the nucleotide, and the Valine (V) is believed to contribute to positioning of the Glycines. The first Glycine and the Valine are invariant in all protein kinases. The second
15 Glycine is almost invariant.

The sequence AIK in the same peptide is also highly conserved and it is located in Domain II, which is also part of the catalytic domain. The Alanine (A) and the Lysine (K) are invariant in all kinases, and the Isoleucine is highly conserved. The Lysine residue appears to be involved in
20 mediating the phosphotransfer reaction (Hawks et al, 1988).

This exon is responsible for the kinase activity of CDC 7. This implies that the CDC 7 coding sequence from the state of the art is not functional.

The novel exon encoded by amino acid sequence SEQ ID No
25 3 (DVIEKKDGPCSGTKGFRAPE) is part of Domain VIII of protein kinases. Mutagenesis has implicated a role of this domain in the catalytic activity (Hawks et al., 1988). In the sequence TKGFRAP, the amino acids Threonine (T), Phenylalanine and Alanine (A) are highly conserved, and the Glutamic Acid (E) is
30 invariant. Moreover, substitution of the corresponding threonine in the yeast CDC7 homologue (position 281 of the yeast CDC7; position 722 in SEQ ID No 1) to a glutamate resulted in a dominant-negative CDC7mutant (Ohtoshi et al. 1997, Mol Gen Genet 254, 562-570).

35 The novel exon, encoded by amino acid sequences SEQ ID No 4 (NIKDIAQLRGSEELWEVAKLHNRESSFPK) is located in Domain XI of protein kinases, and that in the peptide, the first Leucine (L), and the second Lysine (K) are highly conserved and therefore are believed to be quite important for the correct
40 activity of the protein.

In addition, using degenerated CDC27 oligonucleotides, an *Arabidopsis thaliana* cDNA sequence termed CDC27A1 was found, which upon comparison in the above mentioned databanks, showed high homology with an *Arabidopsis thaliana* genomic DNA sequence (accession number AC 001645). Again, the coding sequence of CDC27A1 (SEQ ID NO 9), found by the present inventors, indicated the presence of two additional coding regions in the *Arabidopsis* CDC27, the gene, corresponding with the amino acid sequences given by SEQ ID NOS 6 and 7. Thus, novel DNA replication modulating proteins in plants were found, comprising one or more of the above mentioned novel amino acid sequences.

The novel exon encoded by amino acid sequence SEQ ID No 6 (VNLQLLARCYLSNQAYSAYYILK) is part of a unique NH₂-terminal domain conserved in CDC27 homologues of different origin. The unique domain is located upstream of the NH₂-terminal TPR unit of CDC27 (Tugendreich et al. 1993, Proc Natl Acad Sci USA 90, 10031-10035). The role of this domain is currently not known, but its conservation suggests that it is indispensable for CDC27 function. The NH₂-terminal TPR of CDC27 is not tandemly repeated and spans the amino acid residues 174 to 202 in SEQ ID No 5. Proteins, comprising this novel exon sequence according to the invention may therefore promote APC-substrate action and therewith allowing DNA-replication. On the other hand, a peptide comprising the novel exon sequence may be used to occupy the binding region of the substrates for the APC complex, and therewith inhibiting the complex-substrate interactions, resulting in inactivation of APC and to polyploidization/endoreduplication.

The novel amino acid sequence SEQ ID No 7 (AYMERLILPDELVTEENL) is located just after the last (10th) TPR of CDC27 spanning the amino acid residues 670-703 in SEQ ID No 5. Carboxy-terminal extensions downstream from this 10th TPR and variable in length and sequence are common in all known CDC27 proteins. However, the sequence SEQ ID No 7 shows 50 and 55% homology to the corresponding regions of the CDC27 homologues of *Schizosaccharomyces pombe* and *Aspergillus nidulans*, respectively. Moreover, and previously not recognized, the 25 carboxy-terminal amino acids (ending with SEQ ID No 7) immediately downstream of the 10th TPR compose

aids exists in the SKI3 antiviral protein of *Saccharomyces cerevisiae* (Rhee et al. 1989, Yeast 5, 149-158). Remarkably, three consecutive core amino acids of this TPR, RLI, are also present in SEQ ID No 7 and, although very limited, some further
5 homology can be discovered. Thus, although circumstantial, these data may suggest that SEQ ID No 7 is part of a truncated TPR. If so, the block of tandemly repeated TPRs in CDC27 should be extended from 9 (spanning amino acids 406 to 703 in SEQ ID No 5) to 10 (amino acids 704 to 728 in SEQ ID No 5).
10 Interestingly, it has been suggested that a dimer of the basic 34 amino acid TPR repeat is the more common evolutionary unit (Sikorski et al. 1990, Cell 60, 307-317).

By analyzing patterns of CDC27A1 expression, the present inventors furthermore identified the existence of a second
15 isoform of the CDC27A1 gene. Said isoform, termed CDC27A2 is characterized in that a fragment of 33 nucleotides present in CDC27A1 (nucleotides 1029-1061 of SEQ ID NO 9) is missing in CDC27A2. The nucleotide sequence of the CDC27A2 cDNA is given in SEQ ID NO 14, the corresponding amino acid sequence of the
20 CDC27A2 protein is defined in SEQ ID NO 11. SEQ ID NO 11 is different from SEQ ID NO 5 in that the amino acid sequence 'AIPDVTTLNDP' (SEQ ID NO 12) present in CDC27A1 is absent in CDC27A2.

Another CDC27-like gene from *Arabidopsis thaliana* was
25 identified by the present inventors via in silico cloning. The gene, termed CDC27B has GenBank accession number AC006081 and is annotated as CDC27. However, upon isolation and characterization of the corresponding cDNA, the present inventors noticed that the amino acid sequence predicted and
30 presented in GenBank is lacking the stretch of 161 NH₂-terminal amino acids as given in SEQ ID NO 10.

The cDNA sequence of CDC27B is defined in SEQ ID NO 15 and the derived amino acid sequence of the CDC27B protein is given in SEQ ID NO 13. The full-length CDC27B protein comprises
35 a peptide 75% identical to the peptide as defined in SEQ ID NO 6. As discussed supra, SEQ ID NO 6, and thus also SEQ ID NO 10, are part of a unique NH₂-terminal domain conserved in CDC27 homologues of different origin.

The effect of mutations in one out of the tandem series
40 of TPRs can be very specific. For instance, a point mutation

in the most highly conserved 7th TPR domain of yeast CDC27 results in a greatly reduced affinity for interaction with yeast CDC23, but not for interaction with yeast CDC16 or wild-type CDC27. A single amino acid insertion in the same domain destroys the α -helix and abolishes interaction with wild-type CDC27 as well as CDC16 (Lamb et al. 1994, EMBO J 13, 4321-4328). Moreover, detailed experiments with the human TPR-containing CDC16 and CDC27 homologues and another TPR-containing protein regulating the APC-activity, PP5, revealed that TPR proteins display discriminate binding to other TPR proteins. More specifically for CDC27, deletion of the first TPR domain in this protein abolishes CDC16 binding, but not PP5 binding (Ollendorf and Donoghue 1997, J Biol Chem 272, 32011-32018). Mutagenesis studies with the yeast CDC23 showed that only a few residues in or near the most canonical 6th TPR unit result in temperature-sensitive defects (Sikorski et al. 1993, Mol Cell Biol 13, 1212-1221). Separate TPR domains thus seem to be involved in specific interactions with other proteins and only very limited alterations in these domains seem to be tolerated.

Any erroneous modulation of APC activity, e.g. by mutations in SEQ ID No 6 as part of a conserved sequence in CDC27 proteins and/or SEQ ID No 7 being a putative novel truncated TPR motif in CDC27, will likely result in loss of control over normal DNA replication cycles via the mechanisms described above. Mutations in CDC27 can indeed trigger DNA overreplication and thus the generation of polyploid cells (Heichmann and Roberts 1996, Cell 85, 39-48). Such endoreduplication might be related to cell expansion (Traas et al. 1998, Curr Opin Plant Biol 1, 498-503) and, thus, a higher storage capacity in such polyploid cells. This advantageous property is highly desired in crop plants or parts of plants such as seeds, roots, tubers and fruits.

Modulating the said amino acid sequence would impair the formation of functional APC, whereas cdc27 comprising such a mutation would still be able to interact with the substrate and therewith titrating the substrate out, leading to the abolishment of APC-function in the plant cell, resulting in polyploid cells.

It is to be understood, that DNA replication modulating

proteins according to the present invention, comprising one or more of the above mentioned amino acid sequences, or having 80% amino acid identity therewith, may originate from plant species as well as from other species as long as the said proteins are
5 capable of modulating DNA replication in one or more plant species.

The term "protein" is to be understood as any amino acid sequence having a biological function, optionally modified by e.g. glycosylation. The protein according to the present
10 invention preferably comprises one or more of the amino acid sequences according to c) or d), the respective amino acid identity preferably being at least 50% .

The term "protein" includes single-chain polypeptide molecules as well as multiple-polypeptide complexes where
15 individual constituent polypeptides are linked by covalent or non-covalent means. The term "polypeptide" includes peptides of two or more amino acids in length, typically having more than 5, 10 or 20 amino acids.

20 It will be understood that amino acid sequences of the invention are not limited to the sequences obtained from the particular protein but also include homologous sequences obtained from any source, for example related plant proteins, cellular homologues and synthetic peptides, as well as variants
25 or derivatives thereof.

Thus, the present invention covers variants, homologues or derivatives of the amino acid sequences of the present invention, as well as variants, homologues or derivatives of the nucleotide sequence coding for the amino acid sequences of
30 the present invention.

In the context of the present invention, a homologous sequence is taken to include an amino acid sequence which is at least 50, 60, 70, 80 or 90% identical, preferably at least 95 or 98% identical at the amino acid level over at least 18,
35 preferably all amino acids within the sequences as shown in SEQ ID Nos 2, 3, 4, 6 and 7 in the sequence listing herein. In particular, homology should typically be considered with respect to those regions of the sequence known to be essential for the above discussed functions of the novel amino acid
40 sequences rather than non-essential neighbouring sequences.

Although homology can also be considered in terms of similarity (i.e. amino acid residues having similar chemical properties/functions), in the context of the present invention it is preferred to express homology in terms of sequence identity.

5 Homology comparisons can be conducted by eye, or more usually, with the aid of readily available sequence comparison programs. These commercially available computer programs can calculate % homology between two or more sequences.

10 % Homology may be calculated over contiguous sequences, i.e. one sequence is aligned with the other sequence and each amino acid in one sequence directly compared with the corresponding amino acid in the other sequence, one residue at a time. This is called an "ungapped" alignment. Typically, such ungapped alignments are performed only over a relatively short
15 number of residues (for example less than 50 contiguous amino acids).

Although this is a very simple and consistent method, it fails to take into consideration that, for example, in an otherwise identical pair of sequences, one insertion or
20 deletion will cause the following amino acid residues to be put out of alignment, thus potentially resulting in a large reduction in % homology when a global alignment is performed. Consequently, most sequence comparison methods are designed to produce optimal alignments that take into consideration
25 possible insertions and deletions without penalising unduly the overall homology score. This is achieved by inserting "gaps" in the sequence alignment to try to maximise local homology.

However, these more complex methods assign "gap penalties" to each gap that occurs in the alignment so that,
30 for the same number of identical amino acids, a sequence alignment with as few gaps as possible - reflecting higher relatedness between the two compared sequences - will achieve a higher score than one with many gaps. "Affine gap costs" are typically used that charge a relatively high cost for the
35 existence of a gap and a smaller penalty for each subsequent residue in the gap. This is the most commonly used gap scoring system. High gap penalties will of course produce optimised alignments with fewer gaps. Most alignment programs allow the gap penalties to be modified. However, it is preferred to use
40 the default values when using such software for sequence

comparisons. For example when using the GCG Wisconsin Bestfit package (see below) the default gap penalty for amino acid sequences is -12 for a gap and -4 for each extension.

Calculation of maximum % homology therefore firstly
5 requires the production of an optimal alignment, taking into consideration gap penalties. A suitable computer program for carrying out such an alignment is the GCG Wisconsin Bestfit package (University of Wisconsin, U.S.A.; Devereux et al., 1984, Nucleic Acids Research 12:387). Examples of other
10 software than can perform sequence comparisons include, but are not limited to, the BLAST package (see <http://www.ncbi.nih.gov/BLAST/>), FASTA (Atschul et al., 1990, J. Mol. Biol., 403-410; FASTA is available for online searching at, for example, <http://www.2.ebi.ac.uk.fasta3>) and the
15 GENEWORKS suite of comparison tools. However it is preferred to use the GCG Bestfit program.

Although the final % homology can be measured in terms of identity, the alignment process itself is typically not based on an all-or-nothing pair comparison. Instead, a scaled
20 similarity score matrix is generally used that assigns scores to each pairwise comparison based on chemical similarity or evolutionary distance. An example of such a matrix commonly used is the BLOSUM62 matrix - the default matrix for the BLAST suite of programs. GCG Wisconsin programs generally use either
25 the public default values or a custom symbol comparison table if supplied (see user manual for further details). It is preferred to use the public default values for the GCG package, or in the case of other software, the default matrix, such as BLOSUM62.

30 Once the software has produced an optimal alignment, it is possible to calculate % homology, preferably % sequence identity. The software typically does this as part of the sequence comparison and generates a numerical result.

35 Polypeptide Variants and Derivatives

The terms "variant" or "derivative" in relation to the amino acid sequences of the present invention includes any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) amino acids from or
40 to the sequence providing the resultant amino acid sequence has

similar activity as the polypeptides presented in the sequence listings.

The sequences of the invention may be modified for use in the present invention. Typically, modifications are made that maintain the activity of the sequence. Amino acid substitutions may be made, for example from 1, 2 or 3 to 10, 20 or 30 substitutions provided that the modified sequence retains the relevant activity. E.g. the kinase activity should be maintained in such a variant of a peptide according to the invention comprising SEQ ID NO 2. Amino acid substitutions may include the use of non-naturally occurring analogues, for example to increase blood plasma half-life of a therapeutically administered polypeptide.

Conservative substitutions may be made, for example according to the Table below. Amino acids in the same block in the second column and preferably in the same line in the third column may be substituted for each other:

5	ALIPHATIC	Non-polar	G A P
			I L V
		Polar - uncharged	C S T M
			N Q
	AROMATIC	Polar - charged	D E
			K R
			H F W Y

10 Proteins of the invention are typically made by recombinant means. However they may also be made by synthetic means using techniques well known to skilled persons such as solid phase synthesis. Proteins of the invention may also be produced as fusion proteins, for example to aid in extraction and purification. Examples of fusion protein partners include glutathione-S-transferase (GST), 6xHis, GAL4 (DNA binding and/or transcriptional activation domains) and β -galactosidase. It may also be convenient to include a proteolytic cleavage site between the fusion protein partner and the protein sequence of interest to allow removal of fusion protein sequences. Preferably the fusion protein will not hinder the function of the protein of interest sequence.

Proteins of the invention may be in a substantially isolated form. It will be understood that the protein may be mixed with carriers or diluents which will not interfere with the intended purpose of the protein and still be regarded as substantially isolated. A protein of the invention may also be in a substantially purified form, in which case it will generally comprise the protein in a preparation in which more than 90%, e.g. 95%, 98% or 99% of the protein in the preparation is a protein of the invention.

In a special embodiment, the protein according to the present invention comprises the amino acid sequence as given in SEQ ID NO 1 or NO 5 or NO 11 or NO 13, or has at least 50%, preferably at least 60%, more preferably at least 70, still more preferably 80% and most preferably at least 90% amino acid identity with one of the said sequences. SEQ ID NO 1 relates to the complete amino acid sequence (889 AA) of the novel CDC7 protein according to the present invention comprising SEQ ID

NOS 2, 3 and 4 (AA 411-430, 710-729, 767-795). SEQ ID NO 5 is the complete amino acid sequence (727 AA) of the novel plant CDC27A1 comprising SEQ ID NOS 6 and 7 and 12 (AA 37-60 and AA 711-727 and AA 344-354 respectively). SEQ ID NO 11 is the complete amino acid sequence (716 AA) of the novel plant CDC27A2 comprising SEQ ID NOS 6 and 7 (AA 37-60 and AA 700-716, respectively) but lacking SEQ ID NO 12.

SEQ ID NO 13 is the complete amino acid sequence (739 AA) of the novel plant CDC27B comprising SEQ ID NO 10 (AA-1-161) which itself comprises a peptide 75% identical to SEQ ID NO 6 (AA 36-59).

Although the proteins according to the present invention may be of non-plant origin, as is indicated above, the protein according to the present invention is preferably a plant protein, more preferably a CDC7 or CDC27 protein, or a functional analogue thereof. A functional analogue is to be understood as any protein or peptide having similar biological effects as a plant CDC7 protein or a CDC27 protein, irrespectively of the origin thereof.

Mutein

In another embodiment, the present invention relates to a mutein of the protein according to the present invention, said mutein comprising at least one amino acid substitution, deletion or addition, affecting the DNA replicative effect of the said protein.

As is already indicated above, the proteins according to the present invention are of high interest for an improvement of e.g. agricultural crops or parasite resistance. By substituting, deleting or adding amino acids to the protein according to the present invention, the modulating effect thereof can be affected, which may lead to desirable or improved properties of the protein.

In particular, DNA replication modulating proteins according to the invention may be activated or deions or additions may be situated within or flanking the amino acid sequence, as given by SEQ ID NOS 2, 3, 4, 6, 7, 10 or 12 (or having at least 50% amino acid identity therewith).

DNA replicating modulating proteins according to the invention may also comprise one or more tetratricopeptide

repeat (TPR) domains. Such domains have been identified in CDC27 (amino acid regions 174-202, 403-431, 432-465, 466-499, 500-533, 534-567, 568-601, 602-635, 636-669, 670-703 in SEQ ID No 5; delineation of regions based on the yeast CDC27 homologue; Lamb et al. 1994, EMBO J 13, 4321-4328) as well as in CDC16, CDC23 and many other proteins (Goebel and Yanagida 1991, Trends Biochem Sci 16, 173-177). The function of these TPR domains is to enable the protein to interact with other proteins in the anaphase promoting complex (APC). In the CDC27 protein according to the present invention, a novel TPR or TPR-like domain has been identified which includes SEQ ID No 7. Mutation analysis in TPR domains of yeast CDC27 has revealed that intact TPRs are necessary for CDC27 function (Lamb et al. 1984, EMBO J 13, 4321-4328) and, thus, also for a functional APC. In the absence of CDC27 function, DNA synthesis becomes uncoupled from cell cycle progression resulting in the establishment of polyploid cells (Heichman and Roberts 1996, Cell 85, 39-48).

20 Peptides

Further, the present invention relates to a peptide, comprising

- a) one or more of the amino acid sequences chosen from the group consisting of those given by SEQ ID NOS 2, 3 and 4,
- b) one or more of the amino acid sequences chosen from the group consisting of those, given by SEQ ID NOS 6 and 7,
- c) one or more amino acid sequences having at least 50 % amino acid identity with those of a), or
- d) one or more amino acid sequences having at least 50% amino acid identity with those of b).

These peptides, firstly identified by the present inventors, are or maybe part of important regulatory sites for binding cellular factors or being a substrate for activating/deactivating mechanisms, such as phosphorylation.

Antibodies

Furthermore, the present invention relates to antibodies specifically recognizing a cell cycle interacting protein

according to the invention or parts, i.e. specific fragments or epitopes, of such a protein. The antibodies of the invention can be used to identify and isolate other cell cycle interacting proteins and genes in any organism, preferably plants. These antibodies can be monoclonal antibodies, polyclonal antibodies or synthetic antibodies as well as fragments of antibodies, such as Fab, Fv or scFv fragments etc. Monoclonal antibodies can be prepared, for example, by the techniques as originally described in Köhler and Milstein, Nature 256 (1975), 495, and Galfré, J. Meth. Enzymol. 73 (1981), 3, which comprise the fusion of mouse myeloma cells to spleen cells derived from immunized mammals. Furthermore, antibodies or fragments thereof to the aforementioned peptides can be obtained by using methods which are described, e.g., in Harlow and Lane "Antibodies, A Laboratory Manual", CSH Press, Cold Spring Harbor, 1988. These antibodies can be used, for example, for the immunoprecipitation and immunolocalization of proteins according to the invention as well as for the monitoring of the synthesis of such proteins, for example, in recombinant organisms, and for the identification of compounds interacting with the protein according to the invention. For example, surface plasmon resonance as employed in the BIAcore system can be used to increase the efficiency of phage antibodies selections, yielding a high increment of affinity from a single library of phage antibodies which bind to an epitope of the protein of the invention (Schier, Human Antibodies Hybridomas 7 (1996), 97-105; Malmberg, J. Immunol. Methods 183 (1995), 7-13). In many cases, the binding phenomena of antibodies to antigens is equivalent to other ligand/anti-ligand binding.

DNA sequences

Further, the present invention relates to a non-genomic DNA sequence, coding for a protein or mutin or peptide according to the present invention, or a DNA sequence having a sequence homology of at least 75% with the said sequence, or to the complementary sequence thereof. Also DNA sequences having at least 75% homology with the above mentioned DNA sequences are encompassed within the invention. These sequences are particularly useful in the generation of DNA vectors to

multiply the DNA sequence or to introduce the said sequence in a host organism, in order to obtain the encoded protein. Further said sequences or parts thereof are advantageously used to identify and isolate homologous sequences from other
5 biological species.

The DNA sequence is preferably substantially free of sequences intervening the coding sequence, and is preferably cDNA.

DNA-sequences of the invention comprise nucleic acid
10 sequences encoding the amino acid sequences of the invention. It will be understood by a skilled person that numerous different polynucleotides can encode the same polypeptide as a result of the degeneracy of the genetic code. In addition, it is to be understood that skilled persons may, using routine
15 techniques, make nucleotide substitutions that do not affect the polypeptide sequence encoded by the polynucleotides of the invention to reflect the codon usage of any particular host organism in which the polypeptides of the invention are to be expressed.

20 Polynucleotides of the invention may comprise DNA or RNA. They may be single-stranded or double-stranded. They may also be polynucleotides which include within them synthetic or modified nucleotides. A number of different types of modification to oligonucleotides are known in the art. These
25 include methylphosphonate and phosphorothioate backbones, addition of acridine or polylysine chains at the 3' and/or 5' ends of the molecule. For the purposes of the present invention, it is to be understood that the polynucleotides described herein may be modified by any method available in the
30 art. Such modifications may be carried out in order to enhance the in vivo activity or life span of polynucleotides of the invention.

The terms "variant", "homologue" or "derivative" in relation to the nucleotide sequence of the present invention
35 include any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) nucleic acid from or to the sequence providing the resultant nucleotide sequence codes for a polypeptide, preferably having at least the same activity as sequences presented in the
40 sequence listings.

As indicated above, with respect to sequence homology, preferably there is at least 75%, more preferably at least 85%, more preferably at least 90% homology to the sequences shown in the sequence listing herein. More preferably there is at least 95%, more preferably at least 98%, homology. Nucleotide homology comparisons may be conducted as described above. A preferred sequence comparison program is the GCG Winsconsin Bestfit program described above. The default scoring matrix has a match value of 10 for each identical nucleotide and -9 for each mismatch. The default gap creation penalty is -50 and the default gap extension penalty is -3 for each nucleotide.

The present invention also encompasses nucleotide sequences that are capable of hybridising selectively to the sequences presented herein, or any variant, fragment or derivative thereof, or to the complement of any of the above. Nucleotide sequences are preferably at least 15 nucleotides in length, more preferably at least 20, 30, 40 or 50 nucleotides in length.

The term "hybridization" as used herein shall include "the process by which a strand of nucleic acid joins with a complementary strand through base pairing" as well as the process of amplification as carried out in polymerase chain reaction technologies.

30, for instance at least 40, 60 or 100 or more contiguous nucleotides. Preferred polynucleotides of the invention will comprise regions preferably at least 80 or 90% and more preferably at least 95% homologous to nucleotides (1229-1291), (2126-2187) or (2298-2385) of SEQ ID No 8 or (109-181) or (2125-2181) or (1029-1061) of SEQ ID No 9; or (109-181) or (2092-2148) of SEQ ID NO 14; or (1-483) of SEQ ID NO 15.

Hybridization conditions are based on the melting temperature (T_m) of the nucleic acid binding complex, as taught in Berger and Kimmel (1987, Guide to Molecular Cloning Techniques, Methods in Enzymology, Vol 152, Academic Press, San Diego CA), and confer a defined "stringency" as explained below.

Maximum stringency typically occurs at about $T_m - 5^\circ\text{C}$ (5°C below the T_m of the probe); high stringency at about 5°C to 10°C below T_m ; intermediate stringency at about 10°C to 20°C below T_m ; and low stringency at about 20°C to 25°C below T_m . As will be understood by those of skill in the art, a maximum

stringency hybridization can be used to identify or detect identical polynucleotide sequences while an intermediate (or low) stringency hybridization can be used to identify or detect similar or related polynucleotide sequences.

- 5 In a preferred aspect, the present invention covers nucleotide sequences that can hybridise to the nucleotide sequence of the present invention under stringent conditions (e.g. 65°C and 0.1xSSC {1xSSC = 0.15 M NaCl, 0.015 M Na₃ Citrate pH 7.0}). Where the polynucleotide of the invention is double-stranded,
- 10 both strands of the duplex, either individually or in combination, are encompassed by the present invention. Where the polynucleotide is single-stranded, it is to be understood that the complementary sequence of that polynucleotide is also included within the scope of the present invention.
- 15 Polynucleotides which are not 100% homologous to the sequences of the present invention but fall within the scope of the invention can be obtained in a number of ways. Other variants of the sequences described herein may be obtained for example by probing DNA libraries made from a range of individuals, for
- 20 example individuals from different populations. In addition, other viral/bacterial, or cellular homologues particularly cellular homologues found in plant cells, may be obtained and such homologues and fragments thereof in general will be capable of selectively hybridising to the sequences shown in
- 25 the sequence listing herein. Such sequences may be obtained by probing cDNA libraries made from or genomic DNA libraries from other animal species, and probing such libraries with probes comprising all or part of SEQ ID Nos 8 or 9 or 14 or 15. This may be useful where for example under conditions of medium to
- 30 high stringency. Similar considerations apply to obtaining species homologues and allelic variants of the polypeptide or nucleotide sequences of the invention.
- Variants and strain/species homologues may also be obtained using degenerate PCR which will use primers designed to target
- 35 sequences within the variants and homologues encoding conserved amino acid sequences within the sequences of the present invention. Conserved sequences can be predicted, for example, by aligning the amino acid sequences from several variants/homologues. Sequence alignments can be performed using
- 40 computer software known in the art. For example the GCG

Wisconsin PileUp program is widely used.

The primers used in degenerate PCR will contain one or more degenerate positions and will be used at stringency conditions lower than those used for cloning sequences with single sequence primers against known sequences.

Alternatively, such polynucleotides may be obtained by site directed mutagenesis of characterised sequences, such as SEQ ID No 8 or 9. This may be useful where for example silent codon changes are required to sequences to optimise codon preferences for a particular host cell in which the polynucleotide sequences are being expressed. Other sequence changes may be desired in order to introduce restriction enzyme recognition sites, or to alter the property or function of the polypeptides encoded by the polynucleotides.

Polynucleotides of the invention may be used to produce a primer, e.g. a PCR primer, a primer for an alternative amplification reaction, a probe e.g. labelled with a revealing label by conventional means using radioactive or non-radioactive labels, or the polynucleotides may be cloned into vectors. Such primers, probes and other fragments will be at least 15, preferably at least 20, for example at least 25, 30 or 40 nucleotides in length, and are also encompassed by the term polynucleotides of the invention as used herein.

Polynucleotides such as a DNA polynucleotides and probes according to the invention may be produced recombinantly, synthetically, or by any means available to those of skill in the art. They may also be cloned by standard techniques.

In general, primers will be produced by synthetic means, involving a step wise manufacture of the desired nucleic acid sequence one nucleotide at a time. Techniques for accomplishing this using automated techniques are readily available in the art.

Longer polynucleotides will generally be produced using recombinant means, for example using a PCR (polymerase chain reaction) cloning techniques. This will involve making a pair of primers (e.g. of about 15 to 30 nucleotides) flanking a region of the lipid targeting sequence which it is desired to clone, bringing the primers into contact with mRNA or cDNA obtained from an animal or human cell, performing a polymerase chain reaction under conditions which bring about amplification

of the desired region, isolating the amplified fragment (e.g. by purifying the reaction mixture on an agarose gel) and recovering the amplified DNA. The primers may be designed to contain suitable restriction enzyme recognition sites so that the amplified DNA can be cloned into a suitable cloning vector. For expression of the DNA sequence according to the invention it may in some instances be advantageous to incorporate one or more intervening sequences (introns) in the sequence coding for the protein to be expressed, as in some expression systems, one or more splicing events must take place in order to obtain high expression rates (e.g. for expression of a barley thionin in transgenic tobacco; Carmona et al. 1993, Plant J 3, 457-462). However, in most cases, the coding sequence (i.e. the cDNA), accompanied by the proper regulatory elements, such as promotor and terminator sequences, are sufficient for proper expression.

In a special embodiment (referring to figs 1 and 2), the invention relates to a cDNA sequence, comprising the DNA sequence as given by SEQ ID NO 8 or SEQ ID NO 9 or SEQ ID NO 14 or SEQ ID NO 15, or having a sequence homology with SEQ ID NO 8 or SEQ ID NO 9 or SEQ ID NO 14 or SEQ ID NO 15 of at least 75% or is the complementary sequence thereof. SEQ ID NO 8 is the cDNA sequence of CDC7 of *Arabidopsis thaliana*, comprising the coding sequence for the newly identified amino acid sequences (SEQ ID NOS 2, 3 and 4) as are discussed above. SEQ ID NO 9, is the cDNA sequence of CDC27 of *Arabidopsis thaliana*, includes the sequences coding for the newly identified amino acid sequences (SEQ ID NOS 6 and 7 and 12) as discussed above. SEQ ID NO 14 is the cDNA sequence of CDC27A2 of *Arabidopsis thaliana* and includes the sequences coding for the newly identified amino acid sequences (SEQ ID Nos 6 and 7) as discussed above but lacks the sequence coding for the newly identified amino acid sequence (SEQ ID NO 12). SEQ ID NO 15 is the cDNA sequence of CDC27B of *Arabidopsis thaliana* and includes the sequences coding for the newly identified amino acid sequence (SEQ ID NO 10) as discussed above.

The presence of the amino acid sequences according to the present invention in DNA replication modulating proteins, in particular in CDC7 and CDC27 respectively, may play an

important role in the biological function of the said proteins. Also, the sequences according to SEQ ID NOS 8 and 9 and 14 and 15, or parts thereof, can advantageously be used to isolate and identify homologntary sequence thereof. Such a DNA sequence
5 codes for an amino acid sequence that till now was not known to be part of DNA replication modulating proteins, in particular of CDC7 and CDC27. It was now found, that DNA sequences, corresponding to the nucleotides 1229-1291, 2126-2187 and 2298-2385 of SEQ ID NO 8 code for new amino acid
10 sequences of plant CDC7. The DNA sequence, corresponding to nucleotides 109-181 and 2125-2148 of SEQ ID NO 9 code for novel amino acid sequences of plant CDC27A1, of *Arabidopsis thaliana*. The DNA sequence, corresponding to nucleotides 109-181 and 2092-2148 of SEQ ID NO 14 code for novel amino acid sequences
15 of plant CDC27A2 of *Arabidopsis thaliana*. The DNA sequence, corresponding to nucleotides 1-483 of SEQ ID NO 15 codes for novel amino acid sequence of plant CDC27B of *Arabidopsis thaliana*. Said DNA sequences may therefore in particular be used to identify and isolate genes or gene fragments from other
20 plants or organisms that are homologous to the CDC7 or CDC27 sequence discussed above.

Probes and primers

In a further embodiment, the DNA sequences according to the
25 invention may be used as primers for use in a nucleic acid amplification technique. Said primers can be used in a particular amplification technique to identify and isolate substantially homologous nucleic acid molecules from other plant species. The design and use of said primers is known by
30 the person skilled in the art. Preferably such amplification primers comprise a contiguous sequence of at least 6 nucleotides, in particular 13 nucleotides, preferably 15 to 25 nucleotides or more, identical or complementary to the nucleotide sequence encoding the amino acid sequence of SEQ ID
35 Nos 1-7 and 10-13. Another application is the use as a hybridization probe to identify nucleic acid molecules hybridizing with a nucleic acid molecule of the invention by homology screening of genomic DNA or cDNA libraries. Furthermore, the person skilled in the art is well aware that
40 it is also possible to label such a nucleic acid probe with an

appropriate marker for specific applications, such as for the detection of the presence of a nucleic acid molecule of the invention in a sample derived from an organism, in particular plants. A number of companies such as Pharmacia Biotech
5 (Piscataway NJ), Promega (Madison WI), and US Biochemical Corp (Cleveland OH) supply commercial kits and protocols for these procedures. Suitable reporter molecules or labels include those radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents as well as substrates, cofactors,
10 inhibitors, magnetic particles and the like.

The nucleic acid sequence for a protein of the invention can also be used to generate hybridization probes for mapping the naturally occurring genomic sequence. The sequence may be mapped to a particular chromosome or to a specific region of
15 the chromosome using well known techniques. These include in situ hybridization to chromosomal spreads, flow-sorted chromosomal preparations, or artificial chromosome constructions such as yeast artificial chromosomes, bacterial artificial chromosomes, bacterial P1 constructions or single
20 chromosome cDNA libraries as reviewed in Price (Blood Rev. 7 (1993), 127-134) and Trask (Trends Genet. 7 (1991), 149-154).

Vectors

Polynucleotides of the invention can be incorporated into a
25 recombinant replicable vector. The vector may be used to replicate the nucleic acid in a compatible host cell. Thus in a further embodiment, the invention provides a method of making polynucleotides of the invention by introducing a polynucleotide of the invention into a replicable vector,
30 introducing the vector into a compatible host cell, and growing the host cell under conditions which bring about replication of the vector. The vector may be recovered from the host cell. Suitable host cells include bacteria such as *E. coli*, yeast, mammalian cell lines and other eukaryotic cell lines, for
35 example insect Sf9 cells.

Preferably, a polynucleotide of the invention in a vector is operably linked to a control sequence that is capable of providing for the expression of the coding sequence by the host cell, i.e. the vector is an expression vector. The term
40 "operably linked" means that the components described are in

a relationship permitting them to function in their intended manner. A regulatory sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under condition compatible with the control sequences.

The control sequences may be modified, for example by the addition of further transcriptional regulatory elements to make the level of transcription directed by the control sequences more responsive to transcriptional modulators.

10 Vectors of the invention may be transformed or transfected into a suitable host cell as described below to provide for expression of a protein of the invention. This process may comprise culturing a host cell transformed with an expression vector as described above under conditions to provide for expression by the vector of a coding sequence encoding the protein, and optionally recovering the expressed protein.

The vectors may be for example, plasmid or virus vectors provided with an origin of replication, optionally a promoter for the expression of the said polynucleotide and optionally a regulator of the promoter. The vectors may contain one or more selectable marker genes, for example an ampicillin resistance gene in the case of a bacterial plasmid or a neomycin resistance gene for a mammalian vector. Vectors may be used, for example, to transfect or transform a host cell.

25 Control sequences operably linked to sequences encoding the protein of the invention include promoters/enhancers and other expression regulation signals. These control sequences may be selected to be compatible with the host cell for which the expression vector is designed to be used in. The term promoter is well-known in the art and encompasses nucleic acid regions ranging in size and complexity from minimal promoters to promoters including upstream elements and enhancers.

The promoter is typically selected from promoters which are functional in mammalian cells, although prokaryotic promoters and promoters functional in other eukaryotic cells may be used.

35 The promoter is typically derived from promoter sequences of viral or eukaryotic genes. For example, it may be a promoter derived from the genome of a cell in which expression is to occur. With respect to eukaryotic promoters, they may be promoters that function in a ubiquitous manner (such as

40

promoters of a-actin, b-actin, tubulin) or, alternatively, a tissue-specific manner (such as promoters of the genes for pyruvate kinase). Tissue-specific promoters specific for selected plant tissue cells are particularly preferred, see 5 below in section "transgenic plants".

It may also be advantageous for the promoters to be inducible so that the levels of expression of the heterologous gene can be regulated during the life-time of the cell. Inducible means that the levels of expression obtained using the promoter can 10 be regulated.

In addition, any of these promoters may be modified by the addition of further regulatory sequences, for example enhancer sequences. Chimeric promoters may also be used comprising sequence elements from two or more different promoters 15 described above.

Therefore, the invention relates to DNA vectors, particularly plasmids, cosmids, viruses, bacteriophages and other vectors used conventionally in genetic engineering that comprise a DNA sequence according to the invention. Methods which are well 20 known to those skilled in the art can be used to construct various plasmids and vectors: see for example, the techniques described in Sambrook, Molecular Cloning A Laboratory Manual, Cold Spring Harbor Laboratory (1989) N.Y. and Ausubel, Current Protocols in Molecular Biology, Green Publishing Associates and 25 Wiley Interscience, N.Y. (1989), (1994). Said vector further preferably comprises a promoter, functional in plant cells, operably linked to the DNA sequence, according to the invention. With such a vector, the DNA sequence according to the invention can be expressed in plant cells and may modulate 30 the DNA replication in the said cells.

Identifying derivatives, variants and homologs of the cell cycle interacting proteins of the invention

In another embodiment, the present invention relates to a 35 method for identifying and/or obtaining proteins capable of modulating the DNA replication in plants, comprising a two-hybrid screening assay, using CDC27 or CDC7 polynucleotide sequences as a bait and a cDNA library of a cell suspension culture as prey.

The yeast two-hybrid assay is a genetic strategy developed to identify proteins (encoded by the cDNAs, the 'preys') able to interact in vivo with a known protein (the 'bait'). Interactions between proteins are detected through the reconstitution of the activity of a transcription activator and the subsequent expression of a reporter gene. The cell culture may be from any organism possessing cell cycle interacting proteins such as animals, preferably mammals. Particularly preferred are plant cell suspension cultures such as from *Arabidopsis*. The nucleic acid molecules encoding proteins or peptides identified to interact with CDC7 or CDC27 in the above mentioned assay can be easily obtained and sequenced by methods known in the art. Therefore, the present invention also relates to a DNA sequence encoding a cell cycle interacting protein obtainable by the method of the invention.

Transgenic plants

To analyse the industrial applicabilities of the invention, transformed plants can be made using the nucleotide sequences according to the invention. Such a transformation of the new gene(s), proteins or inactivated variants/mutants thereof will either positively or negatively have an effect on cell division. Methods to modify the expression levels and/or the activity are known to persons skilled in the art and include for instance overexpression, co-suppression, the use of ribozymes, sense and anti-sense strategies, gene silencing approaches. "Sense strand" refers to the strand of a double-stranded DNA molecule that is homologous to a mRNA transcript thereof. The "anti-sense strand" contains an inverted sequence which is complementary to that of the "sense strand".

Hence, the nucleic acid molecules according to the invention are in particular useful for the genetic manipulation of plant cells in order to modify the characteristics of plants and to obtain plants with modified, preferably with improved or useful phenotypes. Similarly, the invention can also be used to modulate the cell division and the growth of cells, preferentially plant cells, in *in vitro* cultures. A transformed plant can thus be obtained by transforming a plant cell with a gene encoding a polypeptide concerned or fragment thereof alone or in combination. For this purpose tissue specific

promoters, in one construct or being present as a separate construct in addition to the sequence concerned, can be used. Thus, the present invention relates to a method for the production of transgenic plants, plant cells or plant tissue
5 comprising the introduction of a nucleic acid molecule or vector of the invention into the genome of said plant, plant cell or plant tissue.

The invention further relates to a method for modulating DNA replication in plant cells, plant parts or plants by conferring
10 to one or more plant cells the capacity to provide a protein, or a mutein thereof according to the invention, in an amount sufficient to modulate DNA replication and/or to block mitosis of the said cells.

In particular, the said capacity is conferred to one or more
15 plant cells, by

a) transforming one or more plant cells with DNA according to the invention or with a vector according to the invention,

b) maintain or culture the plant cells in order to
20 regenerate plant parts or plants from the transformed cells

c) incubating the cells, plant parts or plants at conditions, allowing expression of the DNA according to claim 11 or 12, to produce a protein according to the invention or a mutein thereof according to the invention. For the expression
25 of the nucleic acid molecules according to the invention in sense or antisense orientation in plant cells, the molecules are placed under the control of regulatory elements which ensure the expression in plant cells. These regulatory elements may be heterologous or homologous with respect to the nucleic
30 acid molecule to be expressed as well with respect to the plant species to be transformed. In general, such regulatory elements comprise a promoter active in plant cells. To obtain expression in all tissues of a transgenic plant, preferably constitutive promoters are used, such as the 35 S promoter of CaMV (Odell,
35 Nature 313 (1985), 810-812) or promoters of the polyubiquitin genes of maize (Christensen, Plant Mol. Biol. 18 (1982), 675-689). In order to achieve expression in specific tissues of a transgenic plant it is possible to use tissue specific promoters (see, e.g., Stockhaus, EMBO J. 8 (1989), 2245-2251).
40 Known are also promoters which are specifically active in

tubers of potatoes or in seeds of different plants species, such as maize, Vicia, wheat, barley etc. Inducible promoters may be used in order to be able to exactly control expression. An example for inducible promoters are the promoters of genes
5 encoding heat shock proteins. Also microspore-specific regulatory elements and their uses have been described (WO96/16182). Furthermore, the chemically inducible Tet-system may be employed (Gatz, Mol. Gen. Genet. 227 (1991); 229-237). Further suitable promoters are known to the person skilled in
10 the art and are described, e.g., in Ward (Plant Mol. Biol. 22 (1993), 361-366). The regulatory elements may further comprise transcriptional and/or translational enhancers functional in plants cells. Furthermore, the regulatory elements may include transcription termination signals, such as a poly-A signal,
15 which lead to the addition of a poly A tail to the transcript which may improve its stability.

Methods for the introduction of foreign DNA into plants are also well known in the art. These include, for example, the transformation of plant cells or tissues with T-DNA using
20 *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes*, the fusion of protoplasts, direct gene transfer (see, e.g., EP-A 164 575), injection, electroporation, biolistic methods like particle bombardment, pollen-mediated transformation, plant RNA virus-mediated transformation, liposome-mediated
25 transformation, transformation using wounded or enzyme-degraded immature embryos, or wounded or enzyme-degraded embryogenic callus and other methods known in the art.

In general, the plants which can be modified according to the invention and which either show overexpression of a protein
30 according to the invention or a reduction of the synthesis of such a protein can be derived from any desired plant species. They can be monocotyledonous plants or dicotyledonous plants, preferably they belong to plant species of interest in agriculture, wood culture or horticulture interest, such as
35 crop plants (e.g. maize, rice, barley, wheat, rye, oats etc.), potatoes, oil producing plants (e.g. oilseed rape, sunflower, pea nut, soy bean, etc.), cotton, sugar beet, sugar cane, leguminous plants (e.g. beans, peas etc.), wood producing plants, preferably trees, etc. The invention further relates
40 to progeny of such plants and to plant material such as roots,

flowers, fruit, leaves, pollen, seeds, seedlings or tubers, obtainable from the plant according to the invention.

The invention further relates to a plant cell, transformed with a vector according to the present invention, or comprising DNA according to the present invention. The invention also relates to plants, obtainable by the method according to the present invention and to progeny of such a plant and to plant material, such as roots, flowers, fruit, leaves, pollen, seeds, seedlings or tubers, obtainable from the plant according to the invention.

Mutants

In further embodiments of the invention, expression of dominant negative mutants of CDC7 or CDC27 are used to modulate DNA replication in plant cells, plant tissues, plant organs and/or whole plants. These embodiments involve the overexpression of a mutein or mutant gene according to the present invention which will inhibit the function of a wild-type allele when expressed in the same cell, thereby the phenotype of a transgenic plant, plant organ or plant cell expressing the mutant will be that of a blocked cell cycle progression.

Herskowitz, Nature 329: 219-222 (1987), reviews the inactivation of genes by interference at the protein level, which is achieved through the expression of specific genetic elements encoding a polypeptide comprising both intact, functional domains of the wild type protein as well as nonfunctional domains of the same wild type protein. Such peptides are known as dominant negative mutant proteins. Examples of dominant negative mutants are given below.

30

CDC7 dominant negative mutant - Nematode resistance

In a special embodiment of the present invention, a DNA vector comprises DNA, coding for a mutein according to the present invention, that is operably linked to a nematode-induced promoter, said promoter functional in plant cells. Nematode infection of plants may cause severe problems to plant growth and crop generation. After penetrating the roots of their hosts, nematodes induce, at the infection sites, the development of feeding cells, specialised in the uptake of solutes from the vascular system of the plant. These infection

sites are of crucial importance for the development for the parasite. In this way, root-knot nematodes induce multinucleated giant cells in the infected plant with highly elevated DNA contents. By specifically blocking the DNA synthesis in the feeding cells, the formation of the said multinucleated giant cells may be blocked, so that the nematodes may not further develop. One can contemplate that a CDC7 mutein, which is not further capable to induce the onset of the DNA synthesis, e.g. by loss of one or more phosphorylation sites or loss of binding function to a plant homolog of yeast DBF4 (Jackson et al 1993 Mol Cell Biol 13, 2899-2908) could, when present in sufficient amounts, block the onset of the DNA synthesis. When DNA, coding for such a mutein, and under the control of a promoter, functional in plant cells and inducible by the presence of nematodes in or in the vicinity of the plant cells, is comprised in the plant cells, the mutein can be expressed in the presence or vicinity of nematodes. This may lead to a DNA synthesis block, therewith avoiding further nematode development. The advantage of such a system is the fact that the plant is not producing any heterologous nematocide, that may be harmful for the plant itself. Such a system is not restricted to CDC7. The person, skilled in the art, aware of this application, will be well aware of the possibilities to take other DNA replication modulating proteins, such as CDC27 for developing an analogous anti-nematode system.

CDC27 mutant - Endoreduplication

A further embodiment of the invention involves the down regulation of CDC27. A further embodiment of the invention involves the downregulation of CDC27 resulting in suppression of the APC complex, modulation of DNA replication and/or blocking mitosis. This can be achieved by expression of CDC27 point mutants. An alternative strategy can be envisaged involving a CDC27 mutein consisting of a block of TPR tandem repeats. Such a mutein is still likely to interact with other TPR-containing proteins from the APC such as CDC16 and CDC23 or APC regulator proteins such as PP5. As such, APC component proteins or APC regulator proteins would probably be titrated out and normal APC function be prevented. Based on results

already obtained from experiments designed to delineate TPR domains involved in the interaction between two TPR proteins (Lamb et al. 1984, EMBO J 13, 4321-4328; Ollendorff and Donoghue 1997, J Biol Chem 272, 32011-32018), this strategy might indeed
5 would prove valuable. Overexpression of CDC27 muteins, via the effect on the APC, can be used to enhance endoreduplication in plant cells, plant tissues, plant organs, or whole plants. For example, as is described above, a CDC27 mutein wherein the SEQ ID No 7 has been mutated, leading to the
10 incapability of this mutein to bind with other factors of the APC can be mentioned. The mutated protein would be still able to interact with the substrate, therewith titrating out the APC, abolishing or at least seriously reducing the APC-function, leading to the formation of polyploid cells. Also,
15 mutations in SEQ ID No 6 or 10 could render the mutein incapable of interacting with the substrate but still capable of binding with the other factors of the APC-complex. The result is the generation of a dominant negative, as the complex will not be able to drive the
20 destruction of key components of the cell cycle machinery, responsible to control the number of DNA-replication cycles.

By manipulating the level of endoreduplication one can increase the storage capacity of, for example, endosperm
25 cells. Thus, another aspect of the current invention is that one or more DNA sequences, vectors or proteins, regulatory sequences or recombinant DNA molecules of the invention can be used to modulate, for instance, endoreduplication in storage cells, storage tissues and/or storage organs of
30 plants or parts thereof.

Preferred target storage organs and parts thereof for the modulation of endoreduplication are, for instance, seeds (such as from cereals, als, oilseed crops), roots (such as in sugar beet), tubers (such as in potato) and fruits (such as
35 in vegetables and fruit species). Furthermore it is expected that increased endoreduplication in storage organs and parts thereof correlates with enhanced storage capacity and as such with improved yield. In yet another embodiment of the invention, a plant with modulated endoreduplication in the
40 whole plant or parts thereof can be obtained from a single

plant cell by transforming the cell, in a manner known to the skilled person, with the above-described means.

CDC27 and CDC7 mutants - Sterile plants

5 Another embodiment of the invention relates to a method for modulating DNA replication and the resultant generation of male or female sterile plants. This would be achieved by the expression of dominant negative mutants of either *cdc7* or *cdc27* under the control of very specific promoters - either
10 from male or female gametophytes - to block cell division and disrupt meiosis. The resulting plants would be naturally sterile.

Overexpression of CDC7 and DBF4 activate DNA synthesis

15 Another embodiment of the invention relates to a method for the generation of plant cells, plant tissues, plant organs, or whole plants with the capacity for the overexpression of CDC7 in combination with a plant homolog of Dbf4 thereby modulating DNA replication. Results in yeast indicate that
20 the association of Dbf4 with CDC7 is essential for the G1 to S transition, namely DNA synthesis (Ohtoshi A, Miyake T, Arai K, Masai H; Mol Gen Genet 254(5): 562-70 1997 May 20). Therefore in the present invention, by overexpressing both CDC7 and Dbf4 proteins, one can activate, stimulate or
25 initiate DNA synthesis in cells where DNA synthesis does not normally take place, such as cells that have already gone through the cell cycle. As a consequence the amount of DNA is increased in the cell therewith manipulating the level of endoreduplication as is outlined above.

30

Polyploid plants

Another embodiment of the invention relates to the generation of polyploid plant cells, plant parts or plants.

35

If for example, plant cells are transformed with a vector, comprising the coding sequence of plant CDC27, according to the present invention, under the control of a suitable promoter and optionally other expression controlling
40 elements, these plant cells may produce CDC27. When the said

plant cells produce CDC27 protein in a sufficient amount, extra rounds of DNA replication may take place before mitosis, leading to polyploid cells.

5 Characterisation of CDC7 and CD27 genes

The architecture of the CDC7 and CDC27 genes are illustrated in figures 1 and 2 and 5. Figure 1 illustrates the genomic architecture of the *Arabidopsis* CDC7 gene, wherein the exons are boxed. The numbers above the box indicate the length of the exon, the number below and between two boxes indicates the length of the intron.

The total length of the coding sequence is 2667 nucleotides, coding for 889 amino acids. The fifth, eleventh and thirteenth exons comprise novel coding sequence; in figure 1, the corresponding boxes are black. It is to be understood, and obvious to a skilled person, that the first and the last triplet of the coding sequence of an exon, may partially be encoded by the last two or one nucleotide(s) from the adjacent downstream exon, and, accordingly, by the first two or one nucleotide(s) of the adjacent upstream exon. In figure 2 and 5, the genomic architecture of the CDC27A1 and CDC27B genes, respectively, of *Arabidopsis thaliana* are depicted as explained for figure 1. The second and the sixteenth (last) exon (black in figure 2) comprise novel coding sequences and were not identified in the known genomic CDC27A1 sequence of *Arabidopsis thaliana* (see text). The entire sequence comprises 2184 nucleotides, corresponding to 727 amino acids.

The first 5 exons (black in figure 5) and part of the 6th exon (black in figure 5) comprise novel coding sequences and were not identified in the known genomic CDC27B sequence of *Arabidopsis thaliana* (see text). The entire sequence comprises 2151 nucleotides, corresponding to 716 amino acids.

In figures 3 and 4, the complete cDNA sequence of CDC7 and CDC27A1, respectively, according to the present invention are depicted, with the respective encoded amino acid sequence therebelow. Vertical lines in the nucleotide sequence indicate the exon boundaries, i.e. ²|³ is the boundary between exons 2 and 3. The exon boundaries are

derived from genomic CDC7 and CDC27A1 sequences (see examples 1 and 2 respectively). Such lines are also drawn in the amino acid sequence, although, as is indicated above, the amino acids, flanking such a vertical line, may be partially encoded by the adjacent exon. Exact positioning of the vertical line is in such a case not possible and is set at the left or the right of such an amino acid in an arbitrary manner. See examples 1 and 2 for further details. An alignment of the CDC27A1 (SEQ ID NO 5) and CDC27B (SEQ ID NO 13) amino acid sequences is given in Figure 6 with indication of SEQ ID NOS 6, 7, 10 and 12. Said CDC27A1 and CDC27B sequences are 49% identical when gaps are introduced in the sequences to ensure optimal alignment and maximal identity.

In Figures 7 and 8, the expression of CDC27A and CDC27B genes is illustrated. Figure 7A shows expression of CDC27A genes (both CDC27A1 and CDC27A2 are detected; indicated by the arrows) in several *Arabidopsis thaliana* tissues: 1- etiolated seedlings; 2-flowers; 3-buds; 4-stems; 5-leaves; 6-roots; siliques; - negative control. Figure 7B shows the expression of CDC27A genes in *Arabidopsis thaliana* root cultures treated with different substances: 1-abscisic acid (ABA); 2-2,4-dichlorophenoxyacetic acid (2,4-D); 3-hydroxyurea; 4-kinetin; 5-kinetin + 1-naphthaleneacetic acid (NAA); 6-NAA; 7-oryzalin; 8-starvation; 9-untreated control roots; -negative control. Figure 8A shows the expression of the CDC27B gene in several *Arabidopsis thaliana* tissues as outlined in Figure 7A. Figure 7B illustrates the expression of the CDC27B gene in *Arabidopsis* root cultures treated with different substances as outlined in Figure 7B.

The invention will now be further illustrated by the following examples, that are not intended to limit the scope of the invention.

EXAMPLES

Although in general the techniques mentioned herein are well known in the art, reference may be made in particular to
5 Sambrook et al., Molecular Cloning, A Laboratory Manual (1989) and Ausubel et al., Current Protocols in Molecular Biology (1995), John Wiley & Sons, Inc. Further, scientific explanations and reasonings in the examples are given for illustrative reasons only, without however being bound
10 thereto.

Example 1.

ISOLATION OF AN ARABIDOPSIS CDC7 HOMOLOGUE

15

Conserved regions of the *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe* CDC7 homologue genes were used to synthesize degenerated oligonucleotides to amplify an *Arabidopsis* CDC7 homologue cDNA fragment. These
20 oligonucleotides were as follows:

1 (sense):

5' AAA/G ATA/C/T GGA/C/G/T GAA/G GGA/C/G/T ACA/C/G/T TT
3'

2 (sense):

25 5' ATA/C/T ATA/C/T CAC/T AGA/G GAA/G ATA/C/T AA 3'

3 (antisense)

5' AG C/TTC A/C/G/TGG A/C/G/TGC C/TCT A/GAA A/C/G/TCC 3'

4 (antisense)

5' AC A/C/G/TCC A/C/G/TA/GC A/GCT CCA A/C/G/TAT A/GTC 3'

30

First strand cDNA prepared from whole *Arabidopsis* plants using the Superscript Preamplification System from Life Technologies was used as template in nested PCR
35 reactions. The first reaction was carried using the pair of oligos 1 and 4, and the second reaction used oligos 2 and 3. PCR conditions were essentially as described (Ferreira et al. 1991). A fragment of approximately 650 bp was eluted from an agarose gel, cloned in pGEM-T and sequenced.
40 Sequencing comparison using the GCG-package version 9.1

showed that the deduced amino acid sequence of the PCR fragment has approximately 40% homology to the published yeast CDC7 sequences. This fragment was then used to screen a lambda gt10 cDNA library prepared from total *Arabidopsis* plants. The largest cDNA isolated, approximately 1,2 kb, was completely sequenced by the dideoxy method. This *Arabidopsis* cDNA contains an open reading frame encoded encoding a polypeptide of 384 amino acids (amino acid 473 to amino acid 856 in figure 3). With the SRS search program the EMBL and EMBLnew databanks were screened for gene sequences designated or annotated with the term cdc7. One genomic sequence from *Arabidopsis thaliana* was found (accession number Z97342). This submitted genomic sequence comprised a predicted gene, indicated as "having similarity to protein kinase HSK of fission yeast", having 11 exons and coding for a protein having 829 amino acids.

With the GCG-package version 9.1, the said genomic sequence was compared with the identified partial cDNA sequence, using the "best-fit program". The identified cDNA-sequence covered nucleotides 119827 to 121978 of the genomic sequence of Z97342.

The identified cDNA-sequence did not correspond with the complete coding sequence of the predicted gene on the Z97342 sequence. Within the present cDNA sequence, two additional coding sequences (additional exons) were identified, namely nucleotides no 120770-120709 and 120350-120263 of Z97342, coding for the amino acid sequences of SEQ ID NOS 3 and 4 respectively.

Upon comparison with the genomic *Arabidopsis* sequence, it however appeared that the present cDNA was not complete. To complete our cDNA at the 5' side we used the CAP-finder kit (Clontech), using the primers (CTCTCCCATCTGGTCATGTC, #1; GAACATGCAGTAGCCGTACC, #2) specified for the cDNA, in nested PCR reactions. For the missing 3' end, two nested sequences specific for the cDNA (AAATGGTGCGAACTCAACAC, #2) and (TATGGGAAGTAGCCAAGCTG, #1) and an anchored oligo-dT on the lower strand were used. PCR conditions were essentially as described (Ferreira et al., 1991). The fragments were eluted from agarose gel and cloned using standard techniques and sequenced. The deduced amino acid sequence encoded by

the PCR fragment showed clear homology to the yeast published CDC7 sequences and matched with an the above mentioned *Arabidopsis* genomic sequence. The DNA-fragment, comprising the missing 5' terminal sequence, comprised an additional coding sequence of 63nt (nrs 122340 to 122278 in Z97342) not identified in Z97342, coding for the amino acid sequence of SEQ ID NO 2.

With the obtained sequences, the complete cDNA for the CDC7 homologue of *Arabidopsis thaliana* could be reconstructed, which is illustrated in figure 3 and in SEQ ID NO 8.

The presently identified CDC7 cDNA comprises additional novel coding sequences, corresponding to novel exons (nos 5, 11 and 13 in figure 3), that were not identified in Z97342, and codes for a protein of 890 amino acids.

Example 2. ISOLATION OF THE ARABIDOPSIS CDC27A1 GENE AND cDNA

20

Conserved regions of the published CDC27 homologue genes (Sikorski et al., 1991 Cold Spring Harbor Symposia on Quantitative Biology vol LVI, 663-673, 1991) were used to synthesize degenerated oligonucleotides to amplify *Arabidopsis* CDC27 cDNA. The oligonucleotides were as follows:

1 (sense):

5' TGG GTA/C/G/T TTA/G GCA/C/G/T A/CAA/G GG 3'

2 (sense):

30 5' ATG GAA/C/G/T G/ATT/C/A TA/TC/T AGA/C/G/T AC 3'

3 (antisense)

5' AGA/G CAT/C TAT/C AAT/C GCA/C/G/T TGG 3'

4 (antisense)

5' TA T/A/G AC/T CAT A/C/G/TCC C/TAA A/C/G/CC A/GAA 3'

35 First strand cDNA prepared from flower buds was used as template in nested PCR reactions. The first reaction was carried using the pair of oligos 1 and 4, and the second reaction used oligos 2 and 3. PCR conditions were as described (Ferreira et al., 1991, Plant Cell 3, 531-540), except that the annealing temperature of the first reaction

40

was 45 C, and for the second reaction, 37 C was used. A fragment of approximately 300 bp was eluted from agarose gel and cloned in pGEM-T. Out of 16 clones sequenced, two showed high homology to published CDC27 sequences (Sikorski et al., 1991 Cold Spring Harbor Symposia on Quantitative Biology vol LVI, 663-673, 1991). This fragment was then used to screen a lambda gt10 cDNA library prepared from total *Arabidopsis* plants. The isolated target cDNA, approximately 2,5 kb, was completely sequenced by the dideoxy method and is shown in fig 4 and in SEQ ID nr 9. A combination of restriction enzymes and oligonucleotide subcloning was used to produce the templates for sequencing.

The *Arabidopsis* CDC27A1 cDNA contains one open reading frame, encoding a polypeptide of 727 amino acids (figure 4). With the SRS search program, the databanks EMBL and EMBL new were screened for gene sequences, homologous to the present CDC27 cDNA sequence. A genomic sequence from *Arabidopsis thaliana* (accession number AC001645) was found, comprising 14 exons, coding for a protein of 727 AA. With the GCG-package version 9.1, the present cDNA-sequence was compared with the said genomic *Arabidopsis* sequence (1) using the "best fit"-program. It appeared that the present cDNA comprised additional coding information for two novel exons, namely the second and last exon of the *Arabidopsis* CDC27-gene (exons 2 and 16 in fig 4).

The amino acid sequences encoded by the second and last exon are depicted in SEQ ID NOS 6 and 7 respectively.

Example 3 DOMINANT NEGATIVE MUTANTS OF CDC7

30

Dominant negative mutants of CDC7 (CDC7 DN) are constructed by creating substitution mutations including amino acid residues 1(G), 5(V), 18(A) and 20(K) of SEQ ID No2; amino acid residues 13(T), 16(F), 18(A) and 20(E) of SEQ ID No3; amino acid residues 7(L) and 18(K) of SEQ ID No4.

Substitutions are not conservative. Expression of a CDC7 DN in a whole plant, a plant tissue, a plant organ or a plant cell results in cell cycle arrest at G1/S. These results are in line with the situation in yeast, wherein one such substitution, threonine 13 of SEQ ID No 3 (position 722 in

SEQ ID No 1) to a glutamate has proven to create a dominant

negative CDC7 in yeast. This CDC7 DN is inactive as a kinase but can still bind DBF4, thus preventing activation of wild-type CDC7 molecules (Ohtoshi et al. 1997, Mol Gen Genet 254, 562-570).

The CDC7 DN mutants can be obtained by site-directed mutagenesis using the ExSite PCR-based site-directed mutagenesis kit (Stratagene, La Jolla, CA). Fidelity of the mutagenesis are confirmed by sequencing.

Example 4 MUTANTS OF CDC27

Several types of CDC27 muteins can be considered:

- 15 (1) Insertion of an amino acid such as proline (P) in the amino acid sequence SEQ ID No 7, e.g. behind the tyrosine (Y) residue leads to a loss-of-function of the APC. It is believed that such an insertion deforms the predicted (α -helix of the novel TPR-like domain of which SEQ ID No 7 is part and causes a disturbance of the overall three-dimensional structure of CDC27, therewith titrating out functional proteins of the APC, such as CDC16 or CDC 23, leading to loss of APC function. In line with these results, altering the α -helix structure in one of the TPR units of yeast CDC27 has been proven, and of any of the TPR units has been hypothesized, to destroy CDC27 function (Lamb et al. 1984, EMBO J. 13, 4321-4328).
- 20 (2) Deletion of the NH2-terminal 100 to 220 or 200 to 220 amino acids of CDC27 also leads to loss of function of the APC by titrating out molecules such as APC substrates or APC regulators. This domain encompasses the conserved amino acid sequence SEQ ID No 6 as well as the first TPR unit of CDC27. Deletion of this sequence in human CDC27 abrogates binding of e.g. CDC16, but not of that of e.g. PP5, an APC regulator protein (Ollendorf and Donoghue 1997, J Biol Chem 272, 32011-32018).
- 35 (3) CDC27 muteins consisting of the conserved NH2-terminal domain (containing SEQ ID No6) and 1, 2 or more of the
- 40

downstream TPR units.

- (4) CDC27 muteins consisting of the novel TPR-like domain (ending with SEQ ID No7) preceded by 1, 2 or more of the upstream TPR units.

- 5 Muteins described in (3) and (4) act as those described in (1) or (2).

The point mutants in (1) are obtained by site-directed mutagenesis using the ExSite PCR-based site-directed mutagenesis kit (Stratagene, La Jolla, CA). Fidelity of the
10 mutagenesis are confirmed by sequencing. Deletion mutants in (2), (3) and (4) are obtained by high-fidelity PCR (Expand High Fidelity PCR System, Boehringer, Mannheim) using primers designed to amplify the desired stretches of the CDC27 nucleotide sequence. Primers include extensions
15 recognized by restriction endonucleases to allow easy cloning in a vector such as pUC18. Amplified sequences are checked by nucleotide sequence determination.

Expressing such CDC27 muteins in a whole plant, a plant tissue, a plant organ or a plant cell will cause
20 malfunctioning of the APC and thus repetitive cycles of DNA synthesis without intervening mitosis. This endoreduplication results in a polyploid phenotype.

25 Example 5 NEMATODE RESISTANCE - CDC7 DN

In order to obtain nematode resistance, the CDC7 DN coding sequence is operably linked to a plant promoter responsive to nematode infection and to the NOS polyadenylation site.
30 The ARM1 or Att0728 promoters can be used (Barthels et al. 1997, Plant Cell 9, 2119-2134). The CDC7 DN expression cassette is subsequently transferred to a binary vector such as pGSC1704 and the resulting vector electroporated into *Agrobacterium tumefaciens* C58C1Rif^R (pGV2260). Transformants
35 are selected on streptomycin/spectinomycin containing medium and checked for the presence of the integral transformed binary vector. *Arabidopsis thaliana* Col-0 is transformed with the selected *A. tumefaciens* strain by the floral dip method (Clough and Bent 1998, Plant J 16, 735-743).
40 Transgenic plants are selected after seed germination in the

presence of hygromycin. Selected transgenic lines and untransformed control lines are infected with root knot or cyst nematodes. Successfulness of infection is scored visually two weeks after inoculation (in vitro infection) or
5 six weeks after inoculation (infection of soil-grown plants). Transgenic lines are considered resistant relative to control plants when they display a significant decrease in the number of females or cysts on roots and/or a significantly reduction in nematode feeding sites and/or egg
10 production and/or viable nematodes in the eggs.

Example 6 MALE STERILITY - CDC7 DN and CDC27 muteins

Male sterility in plants are obtained by disrupting normal
15 pollen development. This is achieved by preventing normal cell division of tapetum cells in the anthers. Operably linking CDC7 DN or CDC27 mutein to a tapetum-specific promoter such as Osg6B (Tsuchiya et al. 1995, Plant Cell Physiol 36, 487-494) and to a NOS polyadenylation site will
20 result in a suitable expression cassette. Introduction of this cassette into *A. thaliana* is done as described in example 5. Selected transformant lines have a reduced and/or abnormal pollen formation/development. This is assessed using microscopic methods.

25

Example 7 ENDOREDUPPLICATION - CDC27 muteins

Any of the muteins are operably linked to a constitutive
30 promoter such as the CaMV 35S promoter (Kay et al. 1987, Science 236, 1299-1302) or to a seed endosperm-specific promoter such as from a 2S albumin seed storage protein (Guerche et al. 1990, Plant Cell 2, 469-478) or to the BLZ2 promoter (Carbonero et al, 1999 in press) and to a
35 polyadenylation signal. Such expression cassettes are transferred to *A. thaliana* as described in example 5. Selected transformant lines have a general higher rate of endoreduplicating cells (CaMV 35S promoter) and/or produce seeds with a higher amount of polyploid endosperm cells (2S
40 albumin promoter). Endoreduplication or polyploidism is

assessed in several ways.

- (1) Confocal microscopy is applied to measure the nuclear diameter. Polyploid cells normally have enlarged nuclei in order to harbor the increased DNA content.
- 5 (2) The DNA content of plant cells is measured by flow cytometry (Galbraith et al. 1991, Plant Physiol 96, 985-989).
- (3) The cyclin B-degrading activity of the APC is determined as described by King et al. (1995, Cell 91, 10 279-288).

Example 8 CDC27 GENE EXPRESSION ANALYSIS BY RT-PCR

First-strand cDNA was prepared from RNA isolated from
15 different *Arabidopsis thaliana* tissues (etiolated seedlings, flowers, flower buds; stems; leaves; roots; siliques) and from *Arabidopsis thaliana* root cultures treated for 48 h with different chemical substances (10^{-6} M abscisic acid; 10^{-7} M 2,4-dichlorophenoxyacetic acid; 100 mM hydroxyurea; 10^{-6} M
20 kinetin; 10^{-6} M kinetin + 10^{-6} M 1-naphthaleneacetic acid; 10^{-6} M 1-naphthaleneacetic acid; 2% (w/v) oryzalin). PCR was performed with these cDNAs using CDC27A-specific primers (sense primer 5' CCG TAG TGC TAG AAT AGC A 3' and antisense primer 5' AGT CAG CGT TGA AGT c3') or CDC27B-specific
25 primers (sense primer 5' TCT CTC GAG GAA GAA AGG CAA CAA 3' and antisense primer 5' GGT TCT TGG AGT AGC TAT GGT TTC 3'). The resulting fragments generated by PCR were separated in an agarose gel, blotted to a nylon membrane and hybridized with an 32 P labeled CDC27A or CDC 27B DNA probe. Results are
30 shown in Figure 7 for CDC27A where the arrows indicate the presence of 2 bands, differing by 30 nucleotides. Sequencing of both fragments showed that they are identical, except for the 30 bp insertion. Figure 8 illustrates the results for CDC27B.

35 The pictures in Figures 7 and 8 are representative of 3 independent experiments. Both genes are expressed in all plant tissues, but at reduced levels in open flowers and siliques. Expression of both genes is not drastically affected by hormone treatments, except for a reduction in
40 expression levels observed when roots were incubated with

2,4-D (2,4-dichlorophenoxyacetic acid).

Ubiquitin specific primers were used in separated RT-PCR reactions, using the same first strand cDNAs and, after hybridization, the ubiquitin signals were used to normalize the experiments with CDC27A and CDC27B (data not shown).

While the results of the experiments with hydroxyurea and oryzalin that are shown suggest a reduction in CDC27A expression levels when roots are treated with hydroxyurea.

If these experiments are normalized with the results of

ubiquitin experiments the difference is not significant.

However, a decrease in CDC27B expression is observed in hydroxyurea treated roots, even when the results are normalized with ubiquitin. This result would indicate that CDC27B expression could be cell cycle regulated.

Example 9 ISOLATION OF AN ARABIDOPSIS CDC27A2 cDNA

The RT-PCR products obtained with the CDC27A-specific primers as defined in Example 8 were cloned. CDC27A clones corresponding to the transcripts of different sizes (see Figure 7) were identified and their nucleotide sequences determined. This revealed that both type of CDC27A clones had identical nucleotide sequences with the exception of a stretch of 33 nucleotides which was absent from the shorter CDC27A cDNA. Hence, the longest CDC27A cDNA is referred to as CDC27A1 (SEQ ID NO 9) whereas the shorter CDC27A cDNA is referred to as CDC27A2 (SEQ ID NO 14).

Example 10 ISOLATION OF AN ARABIDOPSIS CDC27B GENE AND cDNA

By means of in silico cloning a second *Arabidopsis thaliana* CDC27 homologue was identified with GenBank accession number AC006081. The GeneMark software was used to predict the exon-intron structure of the gene (see Figure 5) and it was observed that the amino acid sequence of the protein derived from the predicted open reading frame comprised an extra 161 amino acids at the NH₂-terminus as compared to the GenBank sequence. Subsequently the coding region was isolated by PCR on cDNA using primer lying immediately outside of the predicted open reading frame. A product of the expected size

was obtained, cloned and its nucleotide sequence determined to confirm the predicted open reading frame. The primers used to clone the open reading frame were: sense primer 5' TCT CTC GAG GAA GAA AGG CAA CAA 3' and antisense primer 5' GGT TCT TGG AGT AGC TAT GGT TTC 3'. The new Arabidopsis CDC27 homologue is referred to as CDC27B. The CDC27A1 and CDC27B proteins are aligned in Figure 6 and are only 49% identical.

CLAIMS

1. At least partially purified protein, capable of modulating DNA replication in plants, at least comprising in
5 the amino acid sequence
- a) one or more of the amino acid sequences chosen from the group consisting of those, given by SEQ ID NOS 2, 3 and 4,
 - b) one or more of the amino acid sequences chosen from
10 the group consisting of those, given by SEQ ID NOS 6, 7, 10 and 12
 - c) one or more amino acid sequences having at least 50% amino acid identity with those of a), or
 - d) one or more amino acid sequences having at least
15 50% amino acid identity with those of b).
2. Protein according to claim 1, comprising one or more of the amino acid sequences according to c) or d), the respective amino acid identity being at least 90%.
- 20 3. Protein according to claim 1 or 2, having the amino acid sequence as given in SEQ ID NO 1 or NO 5 or NO 11 or NO 13, or having at least 50% amino acid identity with one of the said sequences.
- 25 4. Protein according to one or more of claims 1-3, being a plant CDC7 protein or a functional analogue thereof.
- 30 5. Protein according to one or more claims 1-3, being a plant CDC27 protein or a functional analogue thereof.
- 35 6. Mutein of a protein according to one or more of the preceding claims, comprising at least one amino acid substitution, deletion or addition, affecting the DNA replicative effect of the said protein.
- 40 7. Mutein according to claim 6, wherein at least one of the phosphorylatable amino acids are deleted or substituted by one or more non-phosphorylatable amino acids.

8. Peptide, comprising
- a) one or more of the amino acid sequences chosen from
5 the group consisting of those given by SEQ ID No
2, 3 and 4,
- b) one or more of the amino acid sequences chosen from
the group consisting of those, given by SEQ ID NOS
6, 7, 10 and 12,
- 10 c) one or more amino acid sequences having at least
50% amino acid identity with those of a), or
- d) one or more amino acid sequences having at least
50% amino acid identity with those of b).

15 9. Antibody, specifically recognizing a protein
according to any of the claims 1-5, a mutein according to
any of the claims 6-7 or a peptide according to claim 8.

10. Antibody according to claim 9, being at least
20 partially purified.

11. Non-genomic DNA sequence coding for a protein
according to one or more of claims 1-5, for a mutein
according to claim 6 or 7, or for a peptide according to
25 claim 8, or DNA sequence having a sequence homology of at
least 75% of the said sequence or the complementary DNA
sequence thereof.

12. DNA sequence according to claim 11, being
30 substantially free of sequences intervening the coding
sequence.

13. DNA sequence according to claim 11 or 12,
comprising the DNA sequence as given by SEQ ID no 8 or SEQ
35 ID no 9 or SEQ ID NO 14 or SEQ ID NO 15 or having a sequence
homology with SEQ ID no 8 or SEQ ID no 9 or SEQ ID NO 14 or
SEQ ID NO 15 of at least 75% or the complementary sequence
thereof.

40 14. DNA sequence, coding for a peptide according to

claim 8, corresponding to nucleotides 1229-1291, 2126-2187 or 2298-2385 of SEQ ID No 8, or to nucleotides 109-181 or 2125-2181 or 1029-1061 of SEQ ID No 9, or to nucleotides 109-181 or 2092-2148 of SEQ ID NO 14 or to nucleotides 1-483
5 of SEQ ID NO 15, or a DNA sequence, having a sequence homology of at least 75% to the said sequence or the complementary sequence thereof.

15. DNA vector, at least comprising the DNA sequence
10 according to one of the claims 11-14.

16. DNA vector according to claim 15, further comprising a promoter, functional in plant cells, operably linked to the DNA sequence according to one of the claims
15 11-14.

17. DNA vector according to claim 15 or 16 comprising DNA coding for a mutein according to claim 6 or 7, operably linked to a nematode-induced promoter, functional in plant
20 cells.

18. Method for modulating DNA replication in plant cells, plant parts or plants by conferring to one or more plant cells the capacity to provide a protein according to
25 one or more of claims 1-5, or a mutein thereof according to claim 6 or 7, in an amount sufficient to modulate DNA replication and/or to block mitosis of the said cells.

19. Method according to claim 18, wherein the said
30 capacity is conferred to one or more plant cells, by

- a) transforming one or more plant cells with DNA according to one of the claims 9-12 or with a DNA vector according to one of the claims 13-15,
- b) culturing the plant cells in order to regenerate
35 plant parts or plants from the transformed cells, or
- c) incubating the cells, plant parts or plants at conditions allowing expression of the said DNA to produce the said protein or a mutein.

20. Method according to claim 18 or 19 for the generation of polyploid plant cells, plant parts or plants.

21. Method for identifying and/or obtaining proteins capable of modulating the DNA replication in plants, comprising a two-hybrid screening assay, using CDC27 or CDC7 polynucleotide sequences as a bait and a cDNA library or of a cell suspension culture as a prey.

22. Method for the production of transgenic plants, plant cells or plant tissue, comprising the introduction of a nucleic acid molecule according to any of the claims 11-14 or a vector according to claim 15 or 16 into the genome of said plant, plant cell or plant tissue.

23. Plant cell, transformed with a vector according to one of the claims 15-16, or comprising the DNA according to one of the claims 11-14.

24. Plant, obtainable by the method according to one or more of claims 18-19.

25. Progeny of a plant according to claim 24.

26. Plant material such as roots, flowers, fruit, leaves, pollen, seeds, seedlings or tubers, obtainable from a plant according to claim 24 or 25.

CDC7 Gene Structure




FIGURE 1

CDC27A1 Gene Structure




FIGURE 2

Fig. 3

1 -----¹ATGTCAGAAAACCTCGGAACCGCGTCAACTCGAGAATTCTACA----- 50
TCTAAGTGTCTGTAACGTTACAGTCTTTTGAGCCTTGGCGCAGTTGAGCTCTTAAGATGT
-----¹M S E N S E P R Q L E N S T -
61 -----GCCGGAAGAGAGCTCATTCTCTTAGTCCCACCAATTGAGACGGCAACGACGACCTTAAC----- 120
CGGCTTCTCTCGAGTAAGGAGAATCAGGGTGGTTAAGTCTGCCCTTGCTGCTGGAATTG
A G R E L I P L S P T N S D G N D D L N -
121 -----TATCATCTGCATGCTTTTGAGTTATCTCCTCTCCTACTTTCTTCTGGTCATCCAGAATCT----- 180
ATAGTAGACGTACGAAAACCTCAATAGAGCAGAGGATGAAAGAAGACCAGTAGGTCTTAGA
Y H L H A F E L S R L L L S S G H P E S -
181 -----GTTATAGATCTTTCTTCAAAGTGTACATACTTCCAAGTTTCTCCTAATCTCCTCAAATAT----- 240
CAATATCTAGAAAGAAGTTTACATGTATGAAGGTTCCAAGAGGATTAGAGCAGTTTATA
V I D L S S K C T Y F Q G S P N L V K Y -
241 -----CTTTGCTCGATCCCTAATTCTCCTATTTCCCTTGCCGAAGATGGCTTCACTGTGACTCTC----- 300
GAAACGAGCTAGGGATTAAAGAGGATAAAGGGAACGGCTTCTACCGAAGTGACACTGAGAG
L C S I P N S P I S L A E D G F T V T L -
301 -----TCGCCTGAGTCTCCCTCCGCTCCGGCTAGTTTCGCCTGTAGTTTGGATTGTCAGGAAAAT----- 360
AGCGGACTCAGAGGGAGCGGAGGCGGATCAAAGCGGACATCAAACCTAAACGTCTTTTAA
S P E S P S A P A S F A C S L D L Q E N -
361 -----GTTGTGTTAGAACAGTTTATGGATCCGAGATCTCTCACGCTAAAGCATTGAGAGAGAAT----- 420
CAACACAATCTTGTCAAATACCTAGGCTCTAGAGAGTGCAGATTTCGTAAGCTCTCTCTTA
V V L E Q F M D P R S L T L K H S R E¹₂N -
421 -----GCGGAACAAGAGGAGCTAGAGCTCATGCCCTTGCCCAAAGAAGTCGAAATGATGGAAAC----- 480
CGCCTTGTTCTCCTCGATCTCGAGTACGGTAACGGGTTTTCTTCAGCTTTACTACCTTTG
A E Q E E L E L M P L P K R S R N D G N -

NSDOCID: <WO 0102430A2 1 >

5/17

Q D K S L V Q K T L D Q C K L E G N S K -
ACGTACAGCTGTTCCCTGAGATAAAACACACCAGAAAAAGTAAAGTTATCCAGAAGAGG 1140
TGCATGTCGACAAGGGGACCTCATTTTGTGTGGTCTTTTTCATTTCAATAGGTCTCTCC
T Y S C S P E I K H T R K S K V I Q K R -
AAGCAGAATTTTAAACACCGTTCGTCTTAAAGATCAGAAGGATCAGGCBAAGCATAACACA 1200
TTCGTCTTAAAATTGTGGCAAGCAGAAATTTCTAGTCTTCCAGTCCGTTTCCGTTATTGTGT
K Q N F N T V R L K D Q K | D Q A K E N T -
ATTCCAGATTTTGATTCTTACACTATTGTAGAGGAAGAAGGTTTCAGGTGGCTACGGGATT 1260
TAAGGTCTAAAACCTAAGAATGTGATAACATCTCTTCTTCCAGTCCACCGATGCCCTAA
I P D F D S Y T I V E E E G S G | G V G I -
GTTTATAAGGCAACGAGGAAACTGATGGAACAGAGTTTGCAATTAAATGCCCTCATGTT 1320
CAAATATTCGTTTGTCTCTTTGACTACCTTGTCTCAAACGTTAATTTACGGGAGTACAA
V Y K A T R K T D G T E F A I K | C P H V -
GGCGCTCAGAAGTATTATGTGAATAATGAAATCAGAATGCTGGAGCGTTTGGGGGAAA 1380
CCGCGAGTCTTCATAATACTATTACTTTAGTCTTACGACCTCCGAAACCCCTTT
G A Q K Y Y V N N E I R M L E R F G | G K -
AACTGTATAATAAAGCATGAAGGCTGTCTCAAGAATGGAGATTCTGATTGCATCATCTT 1440
TTGACATATTATTTCGTACTTCCGACAGAGTTCTTACCTCTAAGACTAACGTAGTAGGAA
N C I I K H E G C L K N G D S D C I I L -
GAGCACCTTGAACATGACAGACCTGATTCTTGAAGAGAGAAATAGATGTGTATCAGCTG 1500
CTCGTGGAACCTGTACTGTCTGGACTAAGTAACTTCTCTCTTTTATCTACACATAGTCEAC
E H L E H D R P D S L K R E I D V Y Q L -
CAGTGGTACGGCTACTGCTATGTTCAAAGCTCTATCGAGTCTGCATAAGCAGGGTGTGTT 1560
GTCACCATGCCGATGACGTACAAGTTTCGAGATAGCTCAGACGTATTCTGTCCTCCACACAA
Q W Y G Y C M F K A L S S L H K Q | G V V -
CATAGGGATGTTAAGCCAGGAACTTCTCTCTCTAGGAAGACCAACAAAGGCTATCTC 1620
GTATCCCTACAATTCGGTCTCTTTGAAGGAGAAGAGATCCTTCTGGTGTGTTCCGATAGAG
H R D V K P G N F L F S R K T N K G Y L -
ATTGATTTTAACTTGGCATGATTTGCACCAGAAGTACAGAAGAGCAGATAAATCAAAA

1521 ----- 1580
TAACTAAATTGGAACGGTACCTAAACGGTGGTCTCTGTCCTTCCTGCTATTAGTTTT
I D E N L A M | ⁸ D L H Q K Y R R A D | ⁹ K S K -
GCGGCTTCAGGTCTTCCTACCGCCAGCAAGAAACATCATECATTGGTTAAATCACTCGAT
1681 ----- 1740
CCTCGAAGTCCAGAAGGATGGCGGTCTCTCTGTAGTATGTAACCAATTAGTGAGCTA
A A S G L P T A S K K H E T L V K S L D -
GCGGTAAACCGAGGGACCAACAAACCTTCTCAGAAAACCTTAGCGCCATAAGTTATCAAG
1741 ----- 1800
CGCCATTGGCTCCCTGGTTGTGTTGGAAGAGTCTTTTGAAATCGCGGATTATCATAGTTC
A V N R G T N K P S Q K T L A P N S I K -
AAAGCAGCGGGAAGACAAGAGCTCGGAATGACATGACCAGATGGGAGAGACTCAATAGC
1801 ----- 1860
TTTCGTGCGCCCTTCTGTTCTCGAGCCTTACTGTACTGGTCTACCTCTCTGAGTTATCG
K A A G K T R A R N D M T R W E R L N S -
CAAGGGGCGAAGGGTCTGGCTTAACCTTCAGCTAAAGATGTGACCGCTCAAGGACAAC
1861 ----- 1920
GTTCCCGCTCTTCCCAGACCGAATTGAAGTCGATTTCTACACTGGTCCGTGTTCCCTGTG
Q G A E G S G L T S A K D V T S T R N N -
CCTTCAGGTGAAAAGAGAAGAGAGCCCTTTGCCATGTCATGGAAGAAAAGCGCTTTTAGAT
1921 ----- 1980
GGAAGTCCACTTTCTCTCTCTCGGAAACGGTACAGTACCTTCTTTTCGGGAAAATCTA
P S G E K R R E P L P C H G R K A L L D -
TTTCTGCAAGAGACAATGTCTGTTCCAAFTCCAAACCATGAAGTATCATCCAAAGCTCCT
1981 ----- 2040
AAAGACGTTCTCTGTTACAGACAAGGTTAAGGTTTGGTACTTCATAGTAGGTTTCGAGGA
F L Q E T M S V P I P N H E V S S K A P -
ACGTCTATGAGAAAACGGGTAGCTGCTCTTCCAGGGAAAGCTGAGAAGGAACTTCTTTAT
2041 ----- 2100
TGCAGATACTCTTTTGCCCATCGACGAGAAGGTCCCTTTCCACTCTTCCTTGAAGAAATA
T S M R K R V A A L P G K A E K E L L Y -
CTGACCCCAATGCCACTGTGCTCTAACGGTCCGGCTGAAGCAGGGGACGTAATTGAGAAG
2101 ----- 2160
GACTGGGGTTACGGTGACACGAGATTGCCAGCCGGACTTCGTCCCTGCAATTAATCTTC
L T P M P L C S N G R P E A G | ¹⁰ D V I E K -
AAAGACGGTCCCTGCTCAGGAACCAAGGCTTCCGAGCTCCAGAGTTTGTTCAGATCT
2161 ----- 2220
TTTCTGCCAGGAACGAGTCCCTGGTTTCCGAAGGCTCGAGGTCTCCAAACGAAGTCTAGA
K D G P C S G T K G F R A P E | ¹¹ V C F R S -

2221 TTGCACCAAGGACCTAAGATAGACGTGTGGTCTGCGGGAGTTACCTTGTATACCTCATTA 2280
AACGTGGTTCCTGGATTCTATCTGCACACCGACGCGCTCAATGAAACAATATGGAGTAT
L H Q G E K I D V W S A G V T L L Y L I -
2281 ATGGGAAGGACACCTTTTCACTGGTGACCCCTGAACCTGAACATAAAGGACATGGCACAACTA 2340
TACCTTTCCTGTGGAAAGTGACCACTGGGACTTGTCTTGTATTTCCTGTAAACGTGTGTGAT
M G R T P F T G D B E Q N I K D I A Q L -
2341 CGAGGCAGTGAAGAATTATGGGAAGTAGCCAAAGCTGCACAACCGTGAATCCTCTTTCCCT 2400
GCTCCGTCACTTCTTAATACCTTCATCGGTTCGACGTGTTGGCACTTAGGAGAAAGGCA
R G S E E L W E V A K L E N R E S S E P -
2401 AAGGAATTATACGAGTCAAGGTACTTGAAGGGGATGGAGTTGAGAAAATGGTGCGAACTC 2460
TTCTTAAATATGCTCAGTTCCATGAACCTCCCTACCTCAACTCTTTTACCACGCTTGAG
K E L Y E S R Y L K G M E L R K W C E L -
2461 AACACAAAACGCAGAGAGTTTCTAGACGTAATTCCTACTATCGCTTCTTGACCTCGTTGAT 2520
TTGTGTTTTGCGTCTCTCAAGATCTGCATTAAGGTGATAGCGAAGAACTGGAGCAACTA
N T K R R E F L D V I P L S L L D L V D -
2521 AAATGTTTTGACCGTTAACCCGAGGCGACGAATCAGCGCAGAGGATGCTCTCAAGCAGCAG 2580
TTTACAAACTGGCAATTGGGCTCCGCTGCTTAGTCGCGTCTCTTACGAGAGTTCCGTGCTG
K C L T V N P R R R I S A E D A L K E D -
2581 TTCTTCCATCCAGTACATGAACCCCTTAGAAACCAATGCTCCTTAAACAGCAGCCTACA 2640
AAGAAGGTAGGTCACTGTTACTTTGGGAATCTTTGGTTTACGAGGAATTTGTCGTGCGGTGT
F F H P V H E T L R N Q M L L K Q Q P T -
2641 GTGGTTGCTGACGCAAGTAAGCCAACTCTAAACTATTTACAATTGTAAAGTAAATAAG 2699
CACCAACGACTGCGTCATTCCGTTTGAGATTTGATAAATGTTAACATTTTCATTATTTC
V V A D A V S Q T L N Y L Q L *

1
CCGCTGTAATGTGTGTGTGTCGGAGGCTCCTTGTGTGTGTGTAGCTAACGAGCAGTTAAA 60
.....
1
ATGATGGAGAATCTACTGGCGAATTGTGTCCAGAAAAACCTT 61
----- 120
CCAGTAGTAGTAGTAGTCTACTACCTCTTAGATGACCGCTTAACACAGGTCTTTTTTGGAA
.....
1
M M E N L L A N C V Q K N L
.....
AACCATTTTATGTTTACCAATGCTATCTTCCTTTGCGAACTTCTTCTCGCCCAATTTCCA 121
----- 180
TTGGTAAAATACAAGTGGTTACGATAGAAGGAAACGCTTGAAGAAGAGCGGGTTAAAGGT
N H F M F T N A I F L C E L L L A Q F P
1
TCTGAGGTGAACCTGCAATTGTTAGCCAGGTGTTACTTGAGTAACAGTCAAGCTTATAGT 181
----- 240
AGACTCCACTTGGACGTTAACAATCGGTCCACAATGAACCTCATTTGTCAGTTTCAATATCA
1
S E V N L Q L L A R C Y L S N S Q A Y S
2
GCATATTATATCCTTAAAGTTTCAAAAACGCCTCAGTCTCGGTATTTATTTGCATTCTCA 241
----- 300
CGTATAATATAGGAATTTCCAAGTTTTTTGCGGAGTCAGAGCCATAAATAAACGTAAGAGT
2
A Y Y I L K G S K T P Q S R Y L F A F S
TGCTTTAAGTTGGATCTTCTTGGAGAGGCTGAAGCTGCATTGTTGCCCTGTGAAGATTAT 301
----- 360
ACGAAATTCACCTAGAAGAACCTCTCCGACTTCGACGTAACCAACGGGACACTTCTAATA
C F K L D L L G E A E A A L L P C E D Y
3
GCTGAAGAAAGTTTCTGGTGGTGCAGCTGGGCATTATCTTCTTGGTCTTATATATAGATAT 361
----- 420
CGACTTCTTCAAGGACCACCACGTGACCCGTAATAGAAGAACCAGAATATATATCTATA
3
A E E V P G G A A G H Y L L G L I Y R Y

[illegible]

718
AACTTTTTTAGTGAAGAATTGTCAGCAGAGGCTCAAGAAGAATCTGGGCGCCGCCGCTAGT
901 ----- 960
TTGAAAAAATCACTTCTTAACAGTCGTCCTCCGAGTTCCTTCTTAGACCCGCGGCGGCATCA
718
N F F S E E L S A E | A Q E E S G R R R S
GCTAGAATAGCAGCAAGGAAAAAGAATCCTATGTCGCGAGTCATTGGAAAAGATTCCCAT
961 ----- 1020
CGATCTTATCGTCGTTCCCTTTTCTTAGGATACAGCGTCAGTAAACCTTTTCTAAGGGTA
A R I A A R K K N P M S Q S F G K D S H
TGGTTACATCTTTACCTTCCGAGTCAAACCTATGCACCTTCTCTTTCCCTCGATGATTGGA
1021 ----- 1080
ACCAATGTAGAAAGTGGAAAGGCTCAGTTTGATACGTGGAAGAGAAAGGAGCTACTAACCT
W L H L S P S E S N Y A P S L S S M I G
819
AAATGCAGAATCCAAAGCAGCAAAGAAGCGATTCCCTGATACCGTTACTCTAAATGATCCA
1081 ----- 1140
TTTACGTCTTAGGTTTCGTCGTTTCTTCGCTAAGGACTATGGCAATGAGATTTACTAGGT
819
K C R I Q S S K E A | I P D T V T L N D P
GCAACGACGTCAGGCCAGTCTGTAAAGTGACACTGGAAGCTCTGTTGATGATGAGGAAAAG
1141 ----- 1200
CGTTGCTGCAGTCCGGTCAGACATTCAGTGTGACCTTCGAGACAACCTACTACTCCTTTTC
A T T S G Q S V S D T G S S V D D E E K
TCAAATCCTAGTGAATCTTCCCCGGATCGTTTCAGCCTTATTTCTGGAATTTTCAAGAGTG
1201 ----- 1260
AGTTTAGGATCACTTAGAAGGGGCCTAGCAAAGTCGGAATAAAGACCTTAAAGTCTTCAC
S N P S E S S P D R F S L I S G I S E V
91
CTAGGCATTCTGAAAATTCTTGGAGATGGCCACAGGCATTACATATGTACAAGTGTGAG
1261 ----- 1320
GATCCGTAAGACTTTTAAAGAACCTCTACCGGTGTCCGTAAATGTATACATGTTTACAGTC
91
L G I L K I L G D G H R H L H M Y K C Q

1321 GAAGCTTTGTTGGCATATCAAAAGCTATCTCAGAAACAATACAATACACACTGGGTTCTC 1380
CTTCGAAACAACCGTATAGTTTTTCGATAGAGTCTTTGTTATGTTATGTGTGACCCCAAGAG
1381 E A L L A Y Q K L S Q K Q Y N T H W V L
ATGCAGTTTGGAAAAGCATATTTTGAGCTACAAGACTACTTCAACGCTGACTCTTCCTTT 1440
TACGTCCAACCTTTTCGTATAAAACTCGATGTTCTGATGAAGTTGCGACTGAGAAGGAAA
M Q V G K A Y F E L Q D Y F N A D S S F
1441 ACTCTTGCTCATCAAAAGTATCCTTATGCTTTGGAAGGAATGGATACATACTCCACTGTT 1500
TGAGAACGAGTAGTTTTCATAGGAATACGAAACCTTCCTTACCTATGTATGAGGTGACAA
T L A H Q K Y P Y A L E G M D T Y S T V
1501 CTTTATCACCTGAAAGAAGAGATGAGGTTGGGCTATCTGGCTCAGGAACTGATTTCAGTT 1560
GAAATAGTGGACTTTCTCTCTACTCCAACCCGATAGACCGAGTCTTTGACTAAAGTCAA
L Y H L K E E M R L G Y L A Q E L I S V
1561 GATCGCCTGTCTCCAGAATCCTGTTGTGTCAGTTGGGAACTGTTACAGTTTGCCTAAGGAT 1620
CTAGCGGACAGAGGTCTTAGGACCACACGTCACCCCTTGACAATGTCAAACGCATTCCCTA
D R L S P E S W C A V G N C Y S L R K D
1621 CATGATACTGCTCTCAAAATGTTTCAGAGAGCTATCCAACCTGAATGAAAGATTCAATAT 1680
GTACTATGACGAGAGTTTTACAAAGTCTCTCGATAGGTTGACTTACTTTCTAAGTGTATA
H D T A L K M F Q R A I Q L N E R F T Y
1681 GCACATACCCCTTTGTGGCCACGAGTTTGGCGCATTGGAAGAATTGAGGATGCAGAGAGA 1740
CGTGTATGGGAAACACCGGTGCTCAAACGGCGTAACCTTCTTAAGCTCCTACGTCTCTCT
A H T L C G H E F A A L E E F E D A E R
1741 TGCTACCGGAAGGCTCTGGGCATAGATACGAGACACTATAATGCATGGTACGGTCTTGGA 1800
ACGATGGCCTTCCGAGACCCGTATCTATGCTCTGTGATATTACGTACCATGCCAGAACCT
C Y R K A L G I D T R H Y N A W Y G L G -

DOCID: <WO__0102430A2_I_>

161

2221 TTGATACTACCAGACGAGCTGGTGACGGAGGAAAATTTGTAGATTTTATTGTGCAGGTAAT 2280


AACTATGATGGTCTGCTCGACCACTGCCTCCTTTTAAACATCTAAATAACACGTCCATTA
L I L P D E L V T E E N L *
2281 ACACCAGATTATGTTTTCTCATATAACCCAAAGTCATCTGTAATTTTTCTCATCTTTAGAT 2340

TGTGGTCTAATACAAAGAGTATATTGGGTTTTAGTAGACATTAAAAAGAGTAGAAATCTA
.
2341 CAGTCTTGTGGACTAACCCTAAAACAAAACCTGATTATATAAACTTAGAGGGTAATATTAC 2400

GTCAGAACACCTGATTGGGATTTTGTCTTGACTAATATATTTGAATCTCCCATTTATAATG
.
2401 AGAAAATTGTATAGAGTTGGGTTTGAATTTTCATTTCTTTTCCAAGTTGGAACTTTTGTT 2460

TCTTTTAACATATCTCAACCCAAACTTAAAAGTAAAGAAAAGGTTCAACCTTGAAAACAA
.
2461 CAAA 2512

GTT
.

**FIGURE 5**

SEQ ID NO 6

Cdc27A1 1- MMENILANCVQKNLNHFMTNATFLCELLLAQFPSEVNLQOLLARCYLSNS
Cdc27B 1- MEAMLV-DCVNNSLRHFEVYKNAIFMCERLCAEFSEVNLQOLLATSYLONN

SEQ ID NO 10

SEQ ID NO 6

Cdc27A1 51- QAYSAYYILKGSKTPQSRYLFAFSCFKLDLLGEAEAALLP-CEDYAEVVP
Cdc27B 50- QAYSAYHLLKGTQMAQSRYLFALSCFQMDLLNEAESALCPVNEPGAE-IP

SEQ ID NO 10

Cdc27A1 100- GGAAGHYLLGLIYRYSGRKNCISIQFERMALSFDPWCWEAYGELCSLGAAE
Cdc27B 99- NGAAGHYLLGLIY----KKNAA-QQFKQSLTIDPLLWAAYEELCILGAAE

SEQ ID NO 10

Cdc27A1 150- EASTVFGNVASQRLQKTCVEQRISFSEG-ATIDQ--ITDSKAL--KDTG
Cdc27B 144- EATAVFGETAALSIOKQYMQQ-LSTSLGLNTYNEERNSTSTKNTSSEDYS

SEQ ID NO 10

Cdc27A1 194- LSQTEHIPGENQQDLKIMQQPGDIPPNTDRQLSTNGWDLNTPSPVLLQVM
Cdc27B 193- PROSKHTQSHGLKDISGNFHSBGVNGGV----SNMSFY-NTPSPVAAQLS

Cdc27A1 244- D-APPPLLL NMRRPAV-EG-SLMS-----VHG-VRVRRRRNEFSE---
Cdc27B 238- GIAPPPL-FRNFQ-PAVANPNSLITDSSPKSTVNSTLQAPRRRKFVDEGKL

Cdc27A1 280- -ELSAEAQEEEG-RRRSARIAA-----RKNPMSQSFQKDSHWLHLSP
Cdc27B 286- RKISGRLFSDSGPRR-SSRLSADSGANINSSVATVSGNVNNASKYLGGSK

SEQ ID NO 12

Cdc27A1 321- SESNYAPSLSSMIGKCRIQSSK-----EAIPD-TV-----TLNDPA
Cdc27B 335- -----LSSLALRS-VTLRKGHSWANENM-DEGVRGEPFDDSRPNTAS

Cdc27A1 356- TTSGQSVDSTGSSVDDEEKSNPSE--SSPDRFSLIS-GISEVLSILKILG
Cdc27B 375- TTGSMASND----QEDETMSIGGIAMSSQT----ITIGVSEILNLRTLG

Cdc27A1 403- DGERHLHMYKCOEALLAYOKLSOKQYNTHWVLMQVGKAYFELQDYFNADS
Cdc27B 417- EGCRLSYMYRCOEALDTYMKLPHKHYNTGWVLSQVGKAYFELIDYLEAEK

Cdc27A1 453- SETLAHQKYPYALEGMDTYSTVLYHLKEEMRLGYLAQELISVDRLSPESW
Cdc27B 467- AERLARLASPYCLEGMDIYSTVLYHLKEDMKLSYLAQELISTDRLAPOSW

Cdc27A1 503- CAVGNCYSLRKDHDITALKMFORAIQLNERFTYAHTLCGHEFAALEEFEDA
Cdc27B 517- CAMGNCYSLQKDHETALKNELRAVQLNPRFAYAHTLCGHEYTTLEDFENG

Cdc27A1 553- ERCYRKALGIDTRHYNWYGLGMTYLRQEKFEFAQHOFQALALQINPRSSV
Cdc27B 567- MKSYQNALRVDTTRHYNWYGLGMIYLRQEKLEFSEHHERMAFLINPSSSV

Cdc27A1 603- IMCYYGIALHESKRNDREALMMMEKAVLTDKNPLPKYKKAHILTSLGDYH
Cdc27B 617- IMSYLGTSLHALKRSEALEIMEQAIIVADRNPLPMYOKANILVCLERLD

Cdc27A1 653- KAQKVEELKECAPQESSVHASL-GKIYNOLKQYDKAVLHFGIALDLSPS
Cdc27B 667- EALEVLEELKEYAPSESSVYA-LMGRIYKRRNMHDKAMLHFGIALDMKPP

SEQ ID NO 7

Cdc27A1 702- PSDAVKIKAYMERLILPDELVTRENL
Cdc27B 716- ATDVAAIKAAAMEKLHVPDEIDESP

FIGURE 6




FIGURE 7




FIGURE 8

SEQUENCE LISTING

<110> CropDesign N.V.
Universidade Federal do Rio de Janeiro

<120> Plant DNA replication modulating proteins

<130> A0040067

<140>

<141>

<150> EP99202214.5

<151> 1999-07-05

<160> 15

<170> PatentIn Ver. 2.1

<210> 1

<211> 889

<212> PRT

<213> Arabidopsis thaliana

<400> 1

Met Ser Glu Asn Ser Glu Pro Arg Gln Leu Glu Asn Ser Thr Ala Gly
1 5 10 15

Arg Glu Leu Ile Pro Leu Ser Pro Thr Asn Ser Asp Gly Asn Asp Asp
20 25 30

Leu Asn Tyr His Leu His Ala Phe Glu Leu Ser Arg Leu Leu Leu Ser
35 40 45

Ser Gly His Pro Glu Ser Val Ile Asp Leu Ser Ser Lys Cys Thr Tyr
50 55 60

Phe Gln Gly Ser Pro Asn Leu Val Lys Tyr Leu Cys Ser Ile Pro Asn
65 70 75 80

Ser Pro Ile Ser Leu Ala Glu Asp Gly Phe Thr Val Thr Leu Ser Pro
85 90 95

Glu Ser Pro Ser Ala Pro Ala Ser Phe Ala Cys Ser Leu Asp Leu Gln
100 105 110

Glu Asn Val Val Leu Glu Gln Phe Met Asp Pro Arg Ser Leu Thr Leu
115 120 125

Lys His Ser Arg Glu Asn Ala Glu Gln Glu Glu Leu Glu Leu Met Pro
 130 135 140

Leu Pro Lys Arg Ser Arg Asn Asp Gly Asn Asp Val Asn Tyr Ser Val
 145 150 155 160

Ile Asp Ser Arg Pro Asn Asp Ile Arg Thr Val Ala Cys Gly Thr Met
 165 170 175

Leu Gly Thr Ile Leu Ala Leu Glu Ser Gln Ala Ser Val Phe Asn Leu
 180 185 190

Ser Ala Ser Asn Arg Gly Ile Glu Ala Phe Val Gln Asp His Gln Pro
 195 200 205

Gly Pro Gln Thr Ser Asn Ala Ser Val Asp Val Asn Pro Thr His Arg
 210 215 220

Leu Glu Glu Ser Lys Asn Asp Leu Pro Ser Pro Gln Glu Asp Gly Tyr
 225 230 235 240

Tyr Glu Arg Pro Glu Ile Gly Asp Phe Gln Ile Ala Asp Asn Gln Ile
 245 250 255

Leu Ile Glu Glu Gly Asp Asp Lys Asn Lys Lys Asp Leu Phe Pro Lys
 260 265 270

Gly Glu Ile Gln Thr Asp Ser Val Gln Ser Asp Pro Val Ala Ser Leu
 275 280 285

Met Pro Thr Glu Asn Glu Leu Glu Pro Val Gln Ile Val Asp Asp Thr
 290 295 300

Glu Asp Leu Leu Val Asp Asp His Thr Val Asp Ile Val Ser Thr Pro
 305 310 315 320

Asp Arg Glu Leu Pro Leu Lys Pro Ser Ala Thr Glu Ala Asn Gln Asp
 325 330 335

Lys Ser Leu Val Gln Lys Thr Leu Asp Gln Cys Lys Leu Pro Gly Asn
 340 345 350

Ser Lys Thr Tyr Ser Cys Ser Pro Glu Ile Lys His Thr Arg Lys Ser
 355 360 365

Lys Val Ile Gln Lys Arg Lys Gln Asn Phe Asn Thr Val Arg Leu Lys
 370 375 380

Asp Gln Lys Asp Gln Ala Lys His Asn Thr Ile Pro Asp Phe Asp Ser
 385 390 395 400

Tyr Thr Ile Val Glu Glu Gly Ser Gly Gly Tyr Gly Ile Val Tyr
 405 410 415

Lys Ala Thr Arg Lys Thr Asp Gly Thr Glu Phe Ala Ile Lys Cys Pro
 420 425 430

His Val Gly Ala Gln Lys Tyr Tyr Val Asn Asn Glu Ile Arg Met Leu
 435 440 445

Glu Arg Phe Gly Gly Lys Asn Cys Ile Ile Lys His Glu Gly Cys Leu
 450 455 460

Lys Asn Gly Asp Ser Asp Cys Ile Ile Leu Glu His Leu Glu His Asp
 465 470 475 480

Arg Pro Asp Ser Leu Lys Arg Glu Ile Asp Val Tyr Gln Leu Gln Trp
 485 490 495

Tyr Gly Tyr Cys Met Phe Lys Ala Leu Ser Ser Leu His Lys Gln Gly
 500 505 510

Val Val His Arg Asp Val Lys Pro Gly Asn Phe Leu Phe Ser Arg Lys
 515 520 525

Thr Asn Lys Gly Tyr Leu Ile Asp Phe Asn Leu Ala Met Asp Leu His
 530 535 540

Gln Lys Tyr Arg Arg Ala Asp Lys Ser Lys Ala Ala Ser Gly Leu Pro
 545 550 555 560

Thr Ala Ser Lys Lys His His Thr Leu Val Lys Ser Leu Asp Ala Val
 565 570 575

Asn Arg Gly Thr Asn Lys Pro Ser Gln Lys Thr Leu Ala Pro Asn Ser
 580 585 590

Ile Lys Lys Ala Ala Gly Lys Thr Arg Ala Arg Asn Asp Met Thr Arg
 595 600 605

Trp Glu Arg Leu Asn Ser Gln Gly Ala Glu Gly Ser Gly Leu Thr Ser
 610 615 620

Ala Lys Asp Val Thr Ser Thr Arg Asn Asn Pro Ser Gly Glu Lys Arg
 625 630 635 640

Arg Glu Pro Leu Pro Cys His Gly Arg Lys Ala Leu Leu Asp Phe Leu
 645 650 655

Gln Glu Thr Met Ser Val Pro Ile Pro Asn His Glu Val Ser Ser Lys
 660 665 670

Ala Pro Thr Ser Met Arg Lys Arg Val Ala Ala Leu Pro Gly Lys Ala
 675 680 685

Glu Lys Glu Leu Leu Tyr Leu Thr Pro Met Pro Leu Cys Ser Asn Gly
 690 695 700

Arg Pro Glu Ala Gly Asp Val Ile Glu Lys Lys Asp Gly Pro Cys Ser
 705 710 715 720

Gly Thr Lys Gly Phe Arg Ala Pro Glu Val Cys Phe Arg Ser Leu His
 725 730 735

Gln Gly Pro Lys Ile Asp Val Trp Ser Ala Gly Val Thr Leu Leu Tyr
 740 745 750

Leu Ile Met Gly Arg Thr Pro Phe Thr Gly Asp Pro Glu Gln Asn Ile
 755 760 765

Lys Asp Ile Ala Gln Leu Arg Gly Ser Glu Glu Leu Trp Glu Val Ala
 770 775 780

Lys Leu His Asn Arg Glu Ser Ser Phe Pro Lys Glu Leu Tyr Glu Ser
 785 790 795 800

Arg Tyr Leu Lys Gly Met Glu Leu Arg Lys Trp Cys Glu Leu Asn Thr
 805 810 815

Lys Arg Arg Glu Phe Leu Asp Val Ile Pro Leu Ser Leu Leu Asp Leu
 820 825 830

Val Asp Lys Cys Leu Thr Val Asn Pro Arg Arg Arg Ile Ser Ala Glu
 835 840 845

Asp Ala Leu Lys His Asp Phe Phe His Pro Val His Glu Thr Leu Arg
 850 855 860

Asn Gln Met Leu Leu Lys Gln Gln Pro Thr Val Val Ala Asp Ala Val
 865 870 875 880

Ser Gln Thr Leu Asn Tyr Leu Gln Leu
 885

<210> 2

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 2

Gly Tyr Gly Ile Val Tyr Lys Ala Thr Arg Lys Thr Asp Gly Thr Glu
1 5 10 15

Phe Ala Ile Lys
20

<210> 3

<211> 20

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 3

Asp Val Ile Glu Lys Lys Asp Gly Pro Cys Ser Gly Thr Lys Gly Phe
1 5 10 15

Arg Ala Pro Glu
20

<210> 4

<211> 29

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 4

Asn Ile Lys Asp Ile Ala Gln Leu Arg Gly Ser Glu Glu Leu Trp Glu
1 5 10 15

Val Ala Lys Leu His Asn Arg Glu Ser Ser Phe Pro Lys
20 25

<210> 5

<211> 728

<212> PRT

<213> Arabidopsis thaliana

<400> 5

Met Met Glu Asn Leu Leu Ala Asn Cys Val Gln Lys Asn Leu Asn His
 1 5 10 15

Phe Met Phe Thr Asn Ala Ile Phe Leu Cys Glu Leu Leu Leu Ala Gln
 20 25 30

Phe Pro Ser Glu Val Asn Leu Gln Leu Leu Ala Arg Cys Tyr Leu Ser
 35 40 45

Asn Ser Gln Ala Tyr Ser Ala Tyr Tyr Ile Leu Lys Gly Ser Lys Thr
 50 55 60

Pro Gln Ser Arg Tyr Leu Phe Ala Phe Ser Cys Phe Lys Leu Asp Leu
 65 70 75 80

Leu Gly Glu Ala Glu Ala Ala Leu Leu Pro Cys Glu Asp Tyr Ala Glu
 85 90 95

Glu Val Pro Gly Gly Ala Ala Gly His Tyr Leu Leu Gly Leu Ile Tyr
 100 105 110

Arg Tyr Ser Gly Arg Lys Asn Cys Ser Ile Gln Gln Phe Arg Met Ala
 115 120 125

Leu Ser Phe Asp Pro Leu Cys Trp Glu Ala Tyr Gly Glu Leu Cys Ser
 130 135 140

Leu Gly Ala Ala Glu Glu Ala Ser Thr Val Phe Gly Asn Val Ala Ser
 145 150 155 160

Gln Arg Leu Gln Lys Thr Cys Val Glu Gln Arg Ile Ser Phe Ser Glu
 165 170 175

Gly Ala Thr Ile Asp Gln Ile Thr Asp Ser Asp Lys Ala Leu Lys Asp
 180 185 190

Thr Gly Leu Ser Gln Thr Glu His Ile Pro Gly Glu Asn Gln Gln Asp
 195 200 205

Leu Lys Ile Met Gln Gln Pro Gly Asp Ile Pro Pro Asn Thr Asp Arg

210	215	220
Gln Leu Ser Thr Asn Gly Trp Asp Leu Asn Thr Pro Ser Pro Val Leu		
225	230	235 240
Leu Gln Val Met Asp Ala Leu Pro Pro Leu Leu Leu Lys Asn Met Arg		
245	250	255
Arg Pro Ala Val Glu Gly Ser Leu Met Ser Val His Gly Val Arg Val		
260	265	270
Arg Arg Arg Asn Phe Phe Ser Glu Glu Leu Ser Ala Glu Ala Gln Glu		
275	280	285
Glu Ser Gly Arg Arg Arg Ser Ala Arg Ile Ala Ala Arg Lys Lys Asn		
290	295	300
Pro Met Ser Gln Ser Phe Gly Lys Asp Ser His Trp Leu His Leu Ser		
305	310	315 320
Pro Ser Glu Ser Asn Tyr Ala Pro Ser Leu Ser Ser Met Ile Gly Lys		
325	330	335
Cys Arg Ile Gln Ser Ser Lys Glu Val Ile Pro Asp Thr Val Thr Leu		
340	345	350
Asn Asp Pro Ala Thr Thr Ser Gly Gln Ser Val Ser Asp Ile Gly Ser		
355	360	365
Ser Val Asp Asp Glu Glu Lys Ser Asn Pro Ser Glu Ser Ser Pro Asp		
370	375	380
Arg Phe Ser Leu Ile Ser Gly Ile Ser Glu Val Leu Ser Leu Leu Lys		
385	390	395 400
Ile Leu Gly Asp Gly His Arg His Leu His Met Tyr Lys Cys Gln Glu		
405	410	415
Ala Leu Leu Ala Tyr Gln Lys Leu Ser Gln Lys Gln Tyr Asn Thr His		
420	425	430
Trp Val Leu Met Gln Val Gly Lys Ala Tyr Phe Glu Leu Gln Asp Tyr		
435	440	445
Phe Asn Ala Asp Ser Ser Phe Thr Leu Ala His Gln Lys Tyr Pro Tyr		
450	455	460
Ala Leu Glu Gly Met Asp Thr Tyr Ser Thr Val Leu Tyr His Leu Lys		

465	470	475	480
Glu Glu Met Arg Leu Gly Tyr Leu Ala Gln Glu Leu Ile Ser Val Asp			
485		490	495
Arg Leu Ser Pro Glu Ser Trp Cys Ala Val Gly Asn Cys Tyr Ser Leu			
500	505	510	
Arg Lys Asp His Asp Thr Ala Leu Lys Met Phe Gln Arg Ala Ile Gln			
515	520	525	
Leu Asn Glu Arg Phe Thr Tyr Ala His Thr Leu Cys Gly His Glu Phe			
530	535	540	
Ala Ala Leu Glu Glu Phe Glu Asp Ala Glu Arg Cys Tyr Arg Lys Ala			
545	550	555	560
Leu Gly Ile Asp Thr Arg His Tyr Asn Ala Trp Tyr Gly Leu Gly Met			
565	570	575	
Thr Tyr Leu Arg Gln Glu Lys Phe Glu Phe Ala Gln His Gln Phe Gln			
580	585	590	
Leu Ala Leu Gln Ile Asn Pro Arg Ser Ser Val Ile Met Cys Tyr Tyr			
595	600	605	
Gly Ile Ala Leu His Glu Ser Lys Arg Asn Asp Glu Ala Leu Met Met			
610	615	620	
Met Glu Lys Ala Val Leu Thr Asp Ala Lys Asn Pro Leu Pro Lys Tyr			
625	630	635	640
Tyr Lys Ala His Ile Leu Thr Ser Leu Gly Asp Tyr His Lys Ala Gln			
645	650	655	
Lys Val Leu Glu Glu Leu Lys Glu Cys Ala Pro Gln Glu Ser Ser Val			
660	665	670	
His Ala Ser Leu Gly Lys Ile Tyr Asn Gln Leu Lys Gln Tyr Asp Lys			
675	680	685	
Ala Val Leu His Phe Gly Ile Ala Leu Asp Leu Ser Pro Ser Pro Ser			
690	695	700	
Asp Ala Val Lys Ile Lys Ala Tyr Met Glu Arg Leu Ile Leu Pro Asp			
705	710	715	720
Glu Leu Val Thr Glu Glu Asn Leu			

725

<210> 6

<211> 24

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 6

Val	Asn	Leu	Gln	Leu	Leu	Ala	Arg	Cys	Tyr	Leu	Ser	Asn	Ser	Gln	Ala
1				5					10					15	

Tyr	Ser	Ala	Tyr	Tyr	Ile	Leu	Lys
					20		

<210> 7

<211> 18

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 7

Ala	Tyr	Met	Glu	Arg	Leu	Ile	Leu	Pro	Asp	Glu	Leu	Val	Thr	Glu	Glu
1				5				10						15	

Asn Leu

<210> 8

<211> 2670

<212> DNA

<213> Arabidopsis thaliana

<400> 8

atgtcagaaa	actcggaacc	gcgtcaactc	gagaattcta	cagccggaag	agagctcatt	60
cctcttagtc	ccaccaattc	agacggcaac	gacgacctta	actatcatct	gcatgctttt	120
gagttatctc	gtctcctact	ttcttctggt	catccagaat	ctggtataga	tctttcttca	180
aagtgtacat	acttccaagg	ttctccta	atcttctgctc	gatcccta	240	
tctcctat	cccttgccga	agatggcttc	actgtgactc	tctcgctga	gtctccctcc	300
gctccggcta	gtttcgctg	tagtttgat	ttgcaggaaa	atgttggtt	agaacagttt	360
atggatccga	gatctctcac	gctaaagcat	tcgagagaga	atgcggaaca	agaggagcta	420

```

gagctcatgc cattgcccaa aagaagtcga aatgatggaa acgatgtgaa ttactctgta 480
atagatagca gacctaacga catcagaact gttgcctgtg gaactatgct tgggactatt 540
ttagctcttg aatcccaagc ttcggttttc aatttaagtg catctaaccg aggaatagag 600
gcttttgttc aagatcatca gcctgggccg cagacatcca atgcttcagt ggatgtcaat 660
cctacacatc ggtagagga aagcaagaac gatttgccat ctctcagga ggatggatat 720
tacgagcgac ctgaaattgg agatttccaa attgctgaca accaaatatt aatcgaagaa 780
ggtgatgata aaaataagaa ggatctcttc cctaaggagg agatacaaac tgattctgtg 840
cagtccgac ccgttgccctc attgatgcca acagaaaatg agttagaacc agtgcagatt 900
gtggatgaca ctgaagatct acttgtagat gatcacactg tagacatcgt tagcaccct 960
gacagagagc tgccgttgaa gccttctgct acagaagcta atcaagataa atctttggta 1020
caaaaaactc tggatcaatg caaattgccg ggaaacagca aaacgtacag ctgttcccct 1080
gagataaaac acaccagaaa aagtaaagtt atccagaaga ggaagcagaa ttttaacacc 1140
gttcgtctta aagatcagaa ggatcaggca aagcataaca caattccaga ttttgattct 1200
tacactattg tagaggaaga aggttcagggt ggctacggga ttgtttataa ggcaacgagg 1260
aaaactgatg gaacagagtt tgcaattaaa tgccctcatg ttggcgctca gaagtattat 1320
gtgaataatg aaatcagaat gctggagcgt tttgggggga aaaactgtat aataaagcat 1380
gaaggctgtc tcaagaatgg agattctgat tgcacatcc ttgagcacct tgaacatgac 1440
agacctgatt cattgaagag agaaatagat gtgtatcagc tgcagtggta cggctactgc 1500
atgttcaaaag ctctatcgag tctgcataag cagggtgttg ttcataagga tgtaagcca 1560
ggaaacttcc tcttctctag gaagaccaac aaaggctatc tcattgattt taacctgccc 1620
atggatttgc accagaagta cagaagagca gataaatcaa aagcagcttc aggtcttccc 1680
accgccagca agaaacatca tacattgggt aaatcactcg atgcggtaaa ccgagggacc 1740
aacaacactt ctcagaaaac tttagcgctt aatagtatca agaaagcagc gggaaagaca 1800
agagctcggga atgacatgac cagatgggag agactcaata gccaaagggc agaagggtct 1860
ggcttaactt cagctaaaga tgtgaccagc acaaggaaca acccttcagg tgaaaagaga 1920
agagagcctt tgccatgtca tggaagaaaa gcgcttttag attttctgca agagacaatg 1980
tctgttccaa ttccaaacca tgaagtatca tccaaagctc ctacgtctat gagaaaacgg 2040
gtagctgctc ttccagggaa agctgagaag gaacttcttt atctgacccc aatgccactg 2100
tgctctaacg gtcggcctga agcaggggac gtaattgaga agaaagacgg tccttgctca 2160
ggaaccaaag gcttccgagc tccagaggtt tgcttcagat ctttgcacca aggacctaac 2220
atagacgtgt ggtctgcggg agttactttg ttatacctca taatgggaag gacacctttc 2280
actggtgacc ctgaacagaa cataaaggac attgcacaac tacgaggcag tgaagaatta 2340
tggaagtag ccaagctgca caaccgtgaa tcctctttcc ctaaggaatt atacgagtca 2400
aggtacttga aggggatgga gttgagaaaa tgggtgcgaac tcaacacaaa acgcagagag 2460
tttctagacg taattccact atcgcttctt gacctcgttg ataaatgttt gaccgttaac 2520
ccgaggcgac gaatcagcgc agaggatgct ctcaagcacg acttcttcca tccagtacat 2580
gaaaccctta gaaaccaaat gctccttaaa cagcagccta cagtggttgc tgacgcagta 2640
agccaaactc taaactattt acaattgtaa 2670

```

<210> 9

<211> 2434

<212> DNA

<213> Arabidopsis thaliana

<400> 9

```

atgatggaga atctactggc gaattgtgtc cagaaaaacc ttaaccattt tatgttcacc 60
aatgctatct tcctttgcga acttcttctc gcccaatttc catctgaggt gaacctgcaa 120

```

```

ttgttagcca ggtgttactt gagtaacagt caagcttata gtgcatatta tacccttaaa 180
ggttcaaaaa cgccctcagtc tcggtattta tttgcattct catgctttaa gttggatctt 240
cttgagagagg ctgaagctgc attggttgccc tgtgaagatt atgctgaaga agttcctggg 300
ggtgcagctg ggcattatct tcttggtctt atatatagat attctgggag gaagaactgt 360
tcaatacaac agtttaggat ggcattgtca tttgatccat tgtgttgga agcatatgga 420
gaactttgta gtttaggtgc cgctgaagaa gcctcaacag ttttcgggaa tgttgcttcc 480
cagcgtctta aaacttgtgt agaacaaaga ataagcttct cagaaggagc aaccatagac 540
cagattacag attctgataa ggccttaaaa gatacagggtt tatcgcaaac agaacacatt 600
ccaggagaga accaacaaga tctgaaaatt atgcagcagc ctggagatat tccaccaaatt 660
actgacaggc aacttagtac aaacggatgg gacttgaaca caccttctcc agtgctttta 720
caggtaatgg atgctccacc gcctctgctt cttagaata tgcgtcgtcc agcagtggaa 780
ggatctttga tgtctgtaca tggagtgcgt gtgcgtcgaa gaaacttttt tagtgaagaa 840
ttgtcagcag aggctcaaga agaactctgg cgccgccgta gtgctagaat agcagcaagg 900
aaaaagaatc ctatgtcgca gtcatttgga aaagattccc attggttaca tctttcacct 960
tccgagtcaa actatgcacc ttctctttcc tcgatgattg gaaaatgcag aatccaaagc 1020
agcaaagaag cgattcctga taccgttact ctaaattgat cagcaacgac gtcaggccag 1080
tctgtaagtg aacttggaag ctctgttgat gatgaggaaa agtcaaattcc tagtgaattct 1140
tccccgagtc gtttcagcct tatttctgga atttcagaag tgctaggcat tctgaaaatt 1200
cttgagagat gccacaggca ttacatatg tacaagtgtc aggaagcttt gttggcatat 1260
caaaagctat ctcagaaaca atacaataca cactgggttc tcatgcagggt tggaaaagca 1320
tattttgagc tacaagacta cttcaacgct gactcttctt ttactcttgc tcatcaaaaag 1380
tacccttatg ctttggaagg aatggataca tactccactg ttctttatca cctgaaagaa 1440
gagatgaggt tgggctatct ggctcaggaa ctgatttcag ttgatcgctt gtctccagaa 1500
tcctggtgtg cagttgggaa ctgttacagt ttgcgtaagg atcatgatac tgctctcaaa 1560
atgtttcaga gagctatcca actgaatgaa agattcacat atgcacatac cctttgtggc 1620
cacgagtttg ccgcatgtga agaattcgag gatgcagaga gatgctaccg gaaggctctg 1680
ggcatagata cgagacacta taatgcatgg tacggtcttg gaatgacctt tcttcgtcag 1740
gagaaattcg agtttgcgca gcatcaattt caactggctc tccaaataaa tccaagatct 1800
tcagtcatca tgtgttacta tgggaattgct ttgcatgagt caaagagaaa cgatgaggcg 1860
ttgatgatga tggagaaggc tgtactcact gatgcaaaga atccgctccc caagtactac 1920
aaggctcaca tattaaccag cctaggtgat tatcaciaag cacagaaagt tttagaagag 1980
ctcaaagaat gtgctcctca agaaagcagt gtccatgcat cgcttggaat aatatacaat 2040
cagctaaagc aatacgacaa agccgtgtta catttcggca ttgctttgga tttaagccct 2100
tctccatctg atgctgtcaa gataaaggct tacatggaga ggttgatact accagacgag 2160
ctggtgacgg aggaaaattt gtagatttat tgtgcaggta atacaccaga ttatgtttct 2220
catataaccc aaagtcatct gtaatttttc tcatctttag atcagtcttg tggactaacc 2280
ctaaaacaaa actgattata taaacttaga gggtaattt acagaaaatt gtatagagtt 2340
gggtttgaat tttcatttct tttccaagtt ggaacttttg ttcaaaaaaa aaaaaaaaaa 2400
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa 2434

```

<210> 10

<211> 161

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 10

Met Glu Ala Met Leu Val Asp Cys Val Asn Asn Ser Leu Arg His Phe
 1 5 10 15

Val Tyr Lys Asn Ala Ile Phe Met Cys Glu Arg Leu Cys Ala Glu Phe
 20 25 30

Pro Ser Glu Val Asn Leu Gln Leu Leu Ala Thr Ser Tyr Leu Gln Asn
 35 40 45

Asn Gln Ala Tyr Ser Ala Tyr His Leu Leu Lys Gly Thr Gln Met Ala
 50 55 60

Gln Ser Arg Tyr Leu Phe Ala Leu Ser Cys Phe Gln Met Asp Leu Leu
 65 70 75 80

Asn Glu Ala Glu Ser Ala Leu Cys Pro Val Asn Glu Pro Gly Ala Glu
 85 90 95

Ile Pro Asn Gly Ala Ala Gly His Tyr Leu Leu Gly Leu Ile Tyr Lys
 100 105 110

Lys Asn Ala Ala Gln Gln Phe Lys Gln Ser Leu Thr Ile Asp Pro Leu
 115 120 125

Leu Trp Ala Ala Tyr Glu Glu Leu Cys Ile Leu Gly Ala Ala Glu Glu
 130 135 140

Ala Thr Ala Val Phe Gly Glu Thr Ala Ala Leu Ser Ile Gln Lys Gln
 145 150 155 160

Tyr

<210> 11

<211> 716

<212> PRT

<213> Arabidopsis thaliana

<400> 11

Met Met Glu Asn Leu Leu Ala Asn Cys Val Gln Lys Asn Leu Asn His
 1 5 10 15

Phe Met Phe Thr Asn Ala Ile Phe Leu Cys Glu Leu Leu Leu Ala Gln
 20 25 30

Phe Pro Ser Glu Val Asn Leu Gln Leu Leu Ala Arg Cys Tyr Leu Ser
 35 40 45

Asn Ser Gln Ala Tyr Ser Ala Tyr Tyr Ile Leu Lys Gly Ser Lys Thr
 50 55 60

Pro Gln Ser Arg Tyr Leu Phe Ala Phe Ser Cys Phe Lys Leu Asp Leu
 65 70 75 80

Leu Gly Glu Ala Glu Ala Ala Leu Leu Pro Cys Glu Asp Tyr Ala Glu
 85 90 95

Glu Val Pro Gly Gly Ala Ala Gly His Tyr Leu Leu Gly Leu Ile Tyr
 100 105 110

Arg Tyr Ser Gly Arg Lys Asn Cys Ser Ile Gln Gln Phe Arg Met Ala
 115 120 125

Leu Ser Phe Asp Pro Leu Cys Trp Glu Ala Tyr Gly Glu Leu Cys Ser
 130 135 140

Leu Gly Ala Ala Glu Glu Ala Ser Thr Val Phe Gly Asn Val Ala Ser
 145 150 155 160

Gln Arg Leu Lys Thr Cys Val Glu Gln Arg Ile Ser Phe Ser Glu Gly
 165 170 175

Ala Thr Ile Asp Gln Ile Thr Asp Ser Asp Lys Ala Leu Lys Asp Thr
 180 185 190

Gly Leu Ser Gln Thr Glu His Ile Pro Gly Glu Asn Gln Gln Asp Leu
 195 200 205

Lys Ile Met Gln Gln Pro Gly Asp Ile Pro Pro Asn Thr Asp Arg Gln
 210 215 220

Leu Ser Thr Asn Gly Trp Asp Leu Asn Thr Pro Ser Pro Val Leu Leu
 225 230 235 240

Gln Val Met Asp Ala Pro Pro Pro Leu Leu Leu Lys Asn Met Arg Arg
 245 250 255

Pro Ala Val Glu Gly Ser Leu Met Ser Val His Gly Val Arg Val Arg
 260 265 270

Arg Arg Asn Phe Phe Ser Glu Glu Leu Ser Ala Glu Ala Gln Glu Glu
 275 280 285

Ser Gly Arg Arg Arg Ser Ala Arg Ile Ala Ala Arg Lys Lys Asn Pro
 290 295 300
 Met Ser Gln Ser Phe Gly Lys Asp Ser His Trp Leu His Leu Ser Pro
 305 310 315 320
 Ser Glu Ser Asn Tyr Ala Pro Ser Leu Ser Ser Met Ile Gly Lys Cys
 325 330 335
 Arg Ile Gln Ser Ser Lys Glu Ala Thr Thr Ser Gly Gln Ser Val Ser
 340 345 350
 Asp Thr Gly Ser Ser Val Asp Asp Glu Glu Lys Ser Asn Pro Ser Glu
 355 360 365
 Ser Ser Pro Asp Arg Phe Ser Leu Ile Ser Gly Ile Ser Glu Val Leu
 370 375 380
 Ser Ile Leu Lys Ile Leu Gly Asp Gly His Arg His Leu His Met Tyr
 385 390 395 400
 Lys Cys Gln Glu Ala Leu Leu Ala Tyr Gln Lys Leu Ser Gln Lys Gln
 405 410 415
 Tyr Asn Thr His Trp Val Leu Met Gln Val Gly Lys Ala Tyr Phe Glu
 420 425 430
 Leu Gln Asp Tyr Phe Asn Ala Asp Ser Ser Phe Thr Leu Ala His Gln
 435 440 445
 Lys Tyr Pro Tyr Ala Leu Glu Gly Met Asp Thr Tyr Ser Thr Val Leu
 450 455 460
 Tyr His Leu Lys Glu Glu Met Arg Leu Gly Tyr Leu Ala Gln Glu Leu
 465 470 475 480
 Ile Ser Val Asp Arg Leu Ser Pro Glu Ser Trp Cys Ala Val Gly Asn
 485 490 495
 Cys Tyr Ser Leu Arg Lys Asp His Asp Thr Ala Leu Lys Met Phe Gln
 500 505 510
 Arg Ala Ile Gln Leu Asn Glu Arg Phe Thr Tyr Ala His Thr Leu Cys
 515 520 525
 Gly His Glu Phe Ala Ala Leu Glu Glu Phe Glu Asp Ala Glu Arg Cys
 530 535 540

Tyr Arg Lys Ala Leu Gly Ile Asp Thr Arg His Tyr Asn Ala Trp Tyr
 545 550 555 560

Gly Leu Gly Met Thr Tyr Leu Arg Gln Glu Lys Phe Glu Phe Ala Gln
 565 570 575

His Gln Phe Gln Leu Ala Leu Gln Ile Asn Pro Arg Ser Ser Val Ile
 580 585 590

Met Cys Tyr Tyr Gly Ile Ala Leu His Glu Ser Lys Arg Asn Asp Glu
 595 600 605

Ala Leu Met Met Met Glu Lys Ala Val Leu Thr Asp Ala Lys Asn Pro
 610 615 620

Leu Pro Lys Tyr Tyr Lys Ala His Ile Leu Thr Ser Leu Gly Asp Tyr
 625 630 635 640

His Lys Ala Gln Lys Val Leu Glu Glu Leu Lys Glu Cys Ala Pro Gln
 645 650 655

Glu Ser Ser Val His Ala Ser Leu Gly Lys Ile Tyr Asn Gln Leu Lys
 660 665 670

Gln Tyr Asp Lys Ala Val Leu His Phe Gly Ile Ala Leu Asp Leu Ser
 675 680 685

Pro Ser Pro Ser Asp Ala Val Lys Ile Lys Ala Tyr Met Glu Arg Leu
 690 695 700

Ile Leu Pro Asp Glu Leu Val Thr Glu Glu Asn Leu
 705 710 715

<210> 12

<211> 12

<212> PRT

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:peptide

<400> 12

Ala Ile Pro Asp Thr Val Thr Leu Asn Asp Pro Ala
 1 5 10

<210> 13

<211> 739

<212> PRT

<213> Arabidopsis thaliana

<400> 13

Met Glu Ala Met Leu Val Asp Cys Val Asn Asn Ser Leu Arg His Phe
 1 5 10 15

Val Tyr Lys Asn Ala Ile Phe Met Cys Glu Arg Leu Cys Ala Glu Phe
 20 25 30

Pro Ser Glu Val Asn Leu Gln Leu Leu Ala Thr Ser Tyr Leu Gln Asn
 35 40 45

Asn Gln Ala Tyr Ser Ala Tyr His Leu Leu Lys Gly Thr Gln Met Ala
 50 55 60

Gln Ser Arg Tyr Leu Phe Ala Leu Ser Cys Phe Gln Met Asp Leu Leu
 65 70 75 80

Asn Glu Ala Glu Ser Ala Leu Cys Pro Val Asn Glu Pro Gly Ala Glu
 85 90 95

Ile Pro Asn Gly Ala Ala Gly His Tyr Leu Leu Gly Leu Ile Tyr Lys
 100 105 110

Lys Asn Ala Ala Gln Gln Phe Lys Gln Ser Leu Thr Ile Asp Pro Leu
 115 120 125

Leu Trp Ala Ala Tyr Glu Glu Leu Cys Ile Leu Gly Ala Ala Glu Glu
 130 135 140

Ala Thr Ala Val Phe Gly Glu Thr Ala Ala Leu Ser Ile Gln Lys Gln
 145 150 155 160

Tyr Met Gln Gln Leu Ser Thr Ser Leu Gly Leu Asn Thr Tyr Asn Glu
 165 170 175

Glu Arg Asn Ser Thr Ser Thr Lys Asn Thr Ser Ser Glu Asp Tyr Ser
 180 185 190

Pro Arg Gln Ser Lys His Thr Gln Ser His Gly Leu Lys Asp Ile Ser
 195 200 205

Gly Asn Phe His Ser His Gly Val Asn Gly Gly Val Ser Asn Met Ser
 210 215 220

Phe Tyr Asn Thr Pro Ser Pro Val Ala Ala Gln Leu Ser Gly Ile Ala

225		230		235		240
Pro Pro Pro Leu Phe Arg Asn Phe Gln Pro Ala Val Ala Asn Pro Asn						
	245		250		255	
Ser Leu Ile Thr Asp Ser Ser Pro Lys Ser Thr Val Asn Ser Thr Leu						
	260		265		270	
Gln Ala Pro Arg Arg Lys Phe Val Asp Glu Gly Lys Leu Arg Lys Ile						
	275		280		285	
Ser Gly Arg Leu Phe Ser Asp Ser Gly Pro Arg Arg Ser Ser Arg Leu						
	290		295		300	
Ser Ala Asp Ser Gly Ala Asn Ile Asn Ser Ser Val Ala Thr Val Ser						
305		310		315		320
Gly Asn Val Asn Asn Ala Ser Lys Tyr Leu Gly Gly Ser Lys Leu Ser						
	325		330		335	
Ser Leu Ala Leu Arg Ser Val Thr Leu Arg Lys Gly His Ser Trp Ala						
	340		345		350	
Asn Glu Asn Met Asp Glu Gly Val Arg Gly Glu Pro Phe Asp Asp Ser						
	355		360		365	
Arg Pro Asn Thr Ala Ser Thr Thr Gly Ser Met Ala Ser Asn Asp Gln						
	370		375		380	
Glu Asp Glu Thr Met Ser Ile Gly Gly Ile Ala Met Ser Ser Gln Thr						
385		390		395		400
Ile Thr Ile Gly Val Ser Glu Ile Leu Asn Leu Leu Arg Thr Leu Gly						
	405		410		415	
Glu Gly Cys Arg Leu Ser Tyr Met Tyr Arg Cys Gln Glu Ala Leu Asp						
	420		425		430	
Thr Tyr Met Lys Leu Pro His Lys His Tyr Asn Thr Gly Trp Val Leu						
	435		440		445	
Ser Gln Val Gly Lys Ala Tyr Phe Glu Leu Ile Asp Tyr Leu Glu Ala						
	450		455		460	
Glu Lys Ala Phe Arg Leu Ala Arg Leu Ala Ser Pro Tyr Cys Leu Glu						
465		470		475		480
Gly Met Asp Ile Tyr Ser Thr Val Leu Tyr His Leu Lys Glu Asp Met						

485

490

495

Lys Leu Ser Tyr Leu Ala Gln Glu Leu Ile Ser Thr Asp Arg Leu Ala
 500 505 510

Pro Gln Ser Trp Cys Ala Met Gly Asn Cys Tyr Ser Leu Gln Lys Asp
 515 520 525

His Glu Thr Ala Leu Lys Asn Phe Leu Arg Ala Val Gln Leu Asn Pro
 530 535 540

Arg Phe Ala Tyr Ala His Thr Leu Cys Gly His Glu Tyr Thr Thr Leu
 545 550 555 560

Glu Asp Phe Glu Asn Gly Met Lys Ser Tyr Gln Asn Ala Leu Arg Val
 565 570 575

Asp Thr Arg His Tyr Asn Ala Trp Tyr Gly Leu Gly Met Ile Tyr Leu
 580 585 590

Arg Gln Glu Lys Leu Glu Phe Ser Glu His His Phe Arg Met Ala Phe
 595 600 605

Leu Ile Asn Pro Ser Ser Ser Val Ile Met Ser Tyr Leu Gly Thr Ser
 610 615 620

Leu His Ala Leu Lys Arg Ser Glu Glu Ala Leu Glu Ile Met Glu Gln
 625 630 635 640

Ala Ile Val Ala Asp Arg Lys Asn Pro Leu Pro Met Tyr Gln Lys Ala
 645 650 655

Asn Ile Leu Val Cys Leu Glu Arg Leu Asp Glu Ala Leu Glu Val Leu
 660 665 670

Glu Glu Leu Lys Glu Tyr Ala Pro Ser Glu Ser Ser Val Tyr Ala Leu
 675 680 685

Met Gly Arg Ile Tyr Lys Arg Arg Asn Met His Asp Lys Ala Met Leu
 690 695 700

His Phe Gly Leu Ala Leu Asp Met Lys Pro Pro Ala Thr Asp Val Ala
 705 710 715 720

Ala Ile Lys Ala Ala Met Glu Lys Leu His Val Pro Asp Glu Ile Asp
 725 730 735

Glu Ser Pro

<210> 14

<211> 2401

<212> DNA

<213> Arabidopsis thaliana

<400> 14

```

atgatggaga atctactggc gaattgtgtc cagaaaaacc ttaaccattt tatgttcacc 60
aatgctatct tcctttgcga acttcttctc gcccaatttc catctgaggt gaacctgcaa 120
ttgttagcca ggtgttactt gagtaacagt caagcttata gtgcatatta tacccttaaa 180
ggttcaaaaa cgcctcagtc tcggtattta tttgcattct catgctttaa gttggatctt 240
cttgagagag ctgaagctgc attgttgccc tgtgaagatt atgctgaaga agttcctggg 300
ggtgcagctg ggcattatct tcttgggtctt atatatagat attctggggag gaagaactgt 360
tcaatacaac agtttaggat ggcattgtca tttgatccat tgtgttggga agcatatgga 420
gaactttgta gtttaggtgc cgctgaagaa gcctcaacag ttttcgggaa tgttgettcc 480
cagcgtctta aaacttgtgt agaacaaaga ataagcttct cagaaggagc aaccatagac 540
cagattacag attctgataa ggccttaaaa gatacagggt tatcgcaaac agaacacatt 600
ccaggagaga accaacaaga tctgaaaatt atgcagcagc ctggagatat tccaccaaatt 660
actgacaggc aacttagtac aaacggatgg gacttgaaca caccttctcc agtgctttta 720
caggtaatgg atgctccacc gcctctgctt cttagaataa tgcgtcgccc agcagtggaa 780
ggatctttga tgtctgtaca tggagtgcgt gtgcgtcgaa gaaacttttt tagtgaagaa 840
ttgtcagcag aggtcaaga agaactctgg cgccgccgta gtgctagaat agcagcaagg 900
aaaaagaatc ctatgtcgca gtcatttggg aaagattccc attggttaca tctttcacct 960
tccgagtcaa actatgcacc ttctctttcc tcgatgattg gaaaatgcag aatccaaagc 1020
agcaaagaag caacgacgtc aggccagtct gtaagtgaac ctggaagctc tgttgatgat 1080
gaggaaaagt caaatcctag tgaatcttcc ccggatcggt tcagccttat ttctggaatt 1140
tcagaagtgc taagcattct gaaaattctt ggagatggcc acaggcattt acatatgtac 1200
aagtgtcagg aagctttgtt ggcatatcaa aagctatctc agaaacaata caatacacac 1260
tgggttctca tgcaggttgg aaaagcatat tttgagctac aagactactt caacgctgac 1320
tcttccttta ctcttgetca tcaaaagtat ccttatgctt tgggaaggaa ggatacatac 1380
tccactgttc tttatcacct gaaagaagag atgaggttgg gctatctggc tcaggaactg 1440
atttcagttg atcgctgtc tccagaatcc tgggtgtgcag ttgggaactg ttacagtttg 1500
cgtaaggatc atgatactgc tctcaaaatg tttcagagag ctatccaact gaatgaaaga 1560
ttcacatatg cacataccct ttgtggccac gagtttgccg cattggaaga attcgaggat 1620
gcagagagat gctaccgga ggctctgggc atagatacga gacactataa tgcattgtac 1680
ggtcttggaa tgacctatct tcgtcaggag aaattcgagt ttgcgcagca tcaatttcaa 1740
ctggctctcc aaataaatcc aagatcttca gtcacatgt gttactatgg aattgctttg 1800
catgagtcaa agagaaacga tgaggcgttg atgatgatgg agaaggctgt actcactgat 1860
gcaaagaatc cgtcccccaa gtactacaag gtcacatat taaccagcct aggtgattat 1920
cacaaagcac agaaagtttt agaagagctc aaagaatgtg ctctcaaga aagcagtgtc 1980
catgcatcgc ttggcaaaat atacaatcag ctaaagcaat acgacaaagc cgtgttacat 2040
ttcggcattg ctttggattt aagcccttct ccatctgatg ctgtcaagat aaaggcttac 2100
atggagaggt tgatactacc agacgagctg gtgacggagg aaaatttcta gatttattgt 2160
gcaggtaata caccagatta tgtttctcat ataaccctaa gtcacatgta atttttctca 2220
tctttagatc agtcttgtgg actaaccta aaacaaaact gattatataa acttagaggg 2280
taatattaca gaaaattgta tagagttggg tttgaatttt catttctttt ccaagttgga 2340

```

acttttgttc aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2400
a 2401

<210> 15

<211> 2220

<212> DNA

<213> *Arabidopsis thaliana*

<400> 15

atggaagcta	tgcttgtgga	ctgtgtaaac	aacagtcttc	gtcattttgt	ctacaaaaat	60
gctattttca	tgtgcgagcg	tctctgcgct	gagtttcctt	ctgagggttaa	tttgcagcta	120
ttagccacca	gctacctgca	gaataatcaa	gcttacagtg	catatcatct	gctaaagggga	180
acacaaatgg	ctcagtccecg	atacttggtc	gcattatcat	gcttccagat	ggaccttctc	240
aatgaagctg	aatctgcact	ctgccctggt	aatgaacctg	gtgcggagat	cccaaaggt	300
gcagcaggcc	attaccttct	tggacttatt	tacaagaaga	atgctgctca	acaatttaaa	360
cagtccttga	caatagaccc	tctactttgg	gctgcatatg	aggaattatg	tatattaggt	420
gctgctgagg	aagcaactgc	agtttttggt	gaaacagctg	ctctctccat	tcaaaagcag	480
tatatgcaac	aactgtcaac	ttccctcggc	ttaaacactt	acaacgagga	acgtaattca	540
acttctacta	aaaacacgag	ttctgaagat	tatagtccaa	ggcagtctaa	acacacacaa	600
agccatggcc	ttaaagatat	ctccggaaat	ttccattctc	atggaggttaa	tggaggtggt	660
tcgaacatgt	cattctataa	tacgccttcg	ccagtggctg	cacagctatc	cggtatagct	720
ccaccaccac	ttttccggaa	ttttcagcca	gctgttgcaa	acccaaactc	ccttattact	780
gacagttctc	caaagtccac	tggttaactct	actcttcaag	cacctagaag	aaagtttgta	840
gatgaaggaa	agttacgtaa	gatttctggc	agactatttt	ctgattctgg	tccacgacgg	900
agttcaagac	tgtctgctga	ttcaggggca	aacattaatt	caagtgttgc	aacagtaagc	960
ggaaatgtga	acaacgcttc	caagtatttg	ggaggttcta	aattgagttc	tttggcactt	1020
cgttctgtaa	cacttcggaa	gggacactcc	tgggcaaag	aaaacatgga	tgaaggggtc	1080
cgtggggaac	cttttgatga	ttcaaggcct	aatactgcct	caacgactgg	ttctatggct	1140
tccaatgatc	aagaagacga	aacaatgtcg	attggtggca	tagcaatgag	ttctcaaaca	1200
atcacaaattg	gtgtttcggg	aatttttaaac	ctccttagga	cactcggaga	agggtgtaga	1260
ctttcataca	tgtacaggtg	tcaggaggca	ctggatacgt	atatgaaact	tccacataag	1320
cattataata	caggatgggt	tctttcccag	gtcgggaaag	catactttga	actaattgac	1380
tatttagagg	ctgaaaaggc	attccgtctt	gcccgctgg	cttctcctta	ttgcttagaa	1440
ggaatggata	tatactctac	ggcctctctat	catttgaagg	aagacatgaa	gctgagttac	1500
ttggctcagg	aactaatatc	aaccgatcgc	ttagctcctc	aatcttggtg	tgctatggga	1560
aattgctata	gcttgcaaaa	ggaccatgag	accgcaactg	agaatttcct	acgagctggt	1620
caactgaatc	caagatttgc	atatgcacat	accttatgtg	gccacgaata	cacaactctt	1680
gaggattttg	agaacggaat	gaaaagttac	caaaacgcac	ttcgtgtaga	tacaagacac	1740
tacaacgcat	ggtacgggct	tggaaatgata	tatctacgcc	aagagaagtt	agagttctca	1800
gagcatcact	tcagaatggc	tttcctaata	aaccgcaggt	cctctgttat	aatgtcttat	1860
ttagggacat	ctttgcatgc	cttgaagaga	agtgaggaag	cactagagat	aatggagcaa	1920
gccatagtag	cagatagaaa	aaacctctt	ccaatgtacc	agaaagctaa	catacttgct	1980
tgcttagaaa	gattagatga	agctctagaa	gttcttgagg	agctcaaaga	gtatgcgcct	2040
tcagagagca	gcgtttacgc	tttaatgggc	aggatctata	agcggcgaaa	catgcacgat	2100
aaagccatgc	ttcatttcgg	tctagcttta	gatatgaaac	cgcttgcaac	tgacgttgct	2160
gcaataaagg	ctgcaatgga	gaaattgcat	gttccagatg	agatcgatga	gagcccgtga	2220

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 January 2001 (11.01.2001)

PCT

(10) International Publication Number
WO 01/02430 A3

(51) International Patent Classification⁷: C12N 15/54,
15/82, 15/29, 9/12, A01H 1/00

(74) Agent: WITTOP KONING, T., H.; Exter Polak & Char-
louis B.V., P.O. Box 3241, NL-2280 GE Rijswijk (NL).

(21) International Application Number: PCT/EP00/06401

(22) International Filing Date: 5 July 2000 (05.07.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
99202214.5 5 July 1999 (05.07.1999) EP

(71) Applicants (for all designated States except US):
CROPDESIGN N.V. [BE/BE]; Technologiepark 3,
B-9052 Gent (BE). UNIVERSIDADE FEDERAL DO
RIO DE JANEIRO [BR/BR]; CEP-21941-590 Rio de
Janeiro, RJ (BR).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

(72) Inventors; and


(75) Inventors/Applicants (for US only): HEMERLY, Adri-
ana, Silva [BR/BR]; Rua Padre Achotegui, 60-1004,
Leblon, CEP-22430-090 Rio de Janeiro, RJ (BR). FER-
REIRA, Paulo, Cavalcanti, Gomes [BR/BR]; Rua Padre
Achotegui, 60/1004, Leblon, CEP-22430-090 Rio de
Janeiro, RJ (BR). ROMBAUTS, Stephane [BE/BE];
Nederpolder 23, B-9000 Gent (BE).

(88) Date of publication of the international search report:
27 September 2001

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: ARABIDOPSIS THALIANA CDC7 AND CDC27 HOMOLOGS

CDC7 Gene Structure



(57) Abstract: The present invention relates to at least partially purified protein, capable of modulating the DNA replication in plants, muteins thereof, DNA coding therefor and to a method to confer to one or more plant cells the capacity to provide such a protein or mutein. The invention also relates to plants, comprising the said DNA and the progeny thereof.

WO 01/02430 A3

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP/00/06401

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/54 C12N15/82 C12N15/29 C12N9/12 A01H1/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N A01H

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	BEVAN, M. ET AL.: "Analysis of 1.9 Mb of contiguous sequence from chromosome 4 of Arabidopsis thaliana" NATURE, vol. 391, no. 6666, 29 January 1998 (1998-01-29), pages 485-488, XP002157232 abstract	14-16
Y	SEQ ID NO: Z97342 and gene no: 4515c page 486; figure 1; table 3 page 487, column 1, line 46 - line 53 -& EMBL database, Heidelberg, FRG Empln accession number Z97342 04 July 1997 BEVAN, M. ET AL.: "Arabidopsis thaliana DNA chromosome 4, ESSA I FCA contig fragment No. 7" -/--	1-4, 6-13, 17-26

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"G" document member of the same patent family

Date of the actual completion of the international search

15 January 2001

Date of mailing of the international search report

18 APR 2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Fuchs, U

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	<p>XP002157234 cited in the application the whole document -& EMBL database, Heidelberg, FRG Trembl accession number 023540 01 January 1998 BEVAN, M. ET AL.: "PROTEIN KINASE HOMOLOG (KINASE LIKE PROTEIN) (Arabidopsis thaliana)" XP002157235 the whole document</p> <p>---</p>	
Y	<p>MASAI, H. ET AL.: "hsk1+, a Schizosaccharomyces pombe gene related to Saccharomyces cerevisiae CDC7, is required for chromosomal replication" EMBO JOURNAL, vol. 14, no. 13, 3 July 1995 (1995-07-03), pages 3094-3104, XP000978701 see especially page 3096; figure 1 the whole document</p> <p>---</p>	<p>1-4, 6-13, 17-26</p>
X	<p>WO 96 31608 A (JOHN INNES CENTRE INNOVATIONS LIMITED) 10 October 1996 (1996-10-10) abstract page 8, line 12 -page 9, line 1 page 10, line 1 - line 20 SEQ ID NOS: 1, 2 and 5 page 12, line 26 -page 21, line 20 page 39 -page 42; claims 3,8-11,13-18 page 45 -page 4; figure 1</p> <p>---</p>	<p>8,11-13, 15,16, 22,23</p>
X	<p>MIZOGUCHI, T. ET AL.: "ATMPKs: a gene family of plant MAP kinases in Arabidopsis thaliana" FEBS LETTERS, vol. 336, no. 3, 28 December 1993 (1993-12-28), pages 440-444, XP002079537 abstract page 441, column 1, line 21 -page 442, column 1, line 9 sequence of ATMPK4 page 441; figure 1</p> <p>---</p>	<p>8,11,12, 15</p>
	<p>---</p> <p>-/--</p>	

INTERNATIONAL SEARCH REPORT

 International Application No
 PC 00/06401

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	OHTOSHI, A. ET AL.: "Analyses of Saccharomyces cerevisiae Cdc7 kinase point mutants: dominant-negative inhibition of DNA replication on overexpression of kinase-negative Cdc7 proteins"	21
A	MOLECULAR AND GENERAL GENETICS, vol. 254, no. 5, 20 May 1997 (1997-05-20), pages 562-570, XP002157233 cited in the application abstract	1-4, 6-20, 22-26
A	page 562, column 2, line 1 -page 563, column 2, line 55 page 564, column 1, line 11 -page 569, column 2, line 59; figures 1-4; tables 1-3 --- HAMER, M.J. ET AL.: "Control of initiation of DNA synthesis in plants" PROCEEDINGS OF THE PHYTOCHEMICAL SOCIETY OF EUROPE, vol. 39, 1996, pages 211-224, XP000926262 page 211, line 1 -page 212, line 20 page 212, line 33 - line 34 page 213, line 32 -page 215, line 11; figure 16.2 page 216, line 37 -page 222, line 16 page 222, line 34 -page 223, line 6 --- WO 98 41642 A (VLAAMS INTERUNIVERSITAIR INSTITUUT VOOR BIOTECHNOLOGIE) 24 September 1998 (1998-09-24) the whole document	1-4,6-26
A	--- HEMERLY, A. ET AL.: "Dominant negative mutants of the Cdc2 kinase uncouple cell division from iterative plant development" EMBO JOURNAL, vol. 14, no. 16, 15 August 1995 (1995-08-15), pages 3925-3936, XP002045514 the whole document -----	1-4,6-26

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 00/06401

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims: it is covered by claims Nos.:

1-3,6-26 partially and 4 completely

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

1. Claims: 1-3, 6-26 partially and 4 completely

At least partially purified protein capable of modulating DNA replication in plants, at least comprising a) one or more of the amino acid sequences of SEQ ID NOS: 2, 3 and 4, b) one or more amino acid sequences having at least 50% amino acid identity with those of a), said protein comprising one or more of the amino acid sequences according to b) having an amino acid identity of at least 90%, said protein having the amino acid sequence as given in SEQ ID NO: 1 or having at least 50% amino acid identity with said sequence, said protein being a plant CDC7 protein or a functional analogue thereof, mutein of said protein, peptide comprising a) one or more of the amino acid sequences of SEQ ID NOS: 2, 3 and 4, b) one or more amino acid sequences having at least 50% amino acid identity with those of a), antibody specifically recognizing said protein, said mutein or said peptide, non-genomic DNA sequence coding for said protein, said mutein or said peptide or DNA sequence having a sequence homology of at least 75% of said sequence or complementary DNA sequence thereof, said DNA sequence comprising the DNA sequence as given by SEQ ID NO: 8 or having a sequence homology with said sequence of at least 75% or complementary sequence thereof, DNA sequence coding for said peptide corresponding to nucleotides 1229-1291, 2126-2187 or 2298-2385 of SEQ ID NO: 8 or a DNA sequence having a sequence homology of at least 75% to said sequence or complementary sequence thereof, DNA vector comprising said DNA sequence, method for modulating DNA replication in plant cells, plant parts or plants by conferring the capacity to provide said protein or said mutein in an amount sufficient to modulate DNA replication and/or block mitosis of said cells, method for identifying and/or obtaining proteins capable of modulating the DNA replication in plants comprising a two-hybrid screening assay using CDC7 polynucleotide sequences as bait and a cDNA library of a cell suspension culture as prey, method for the production of transgenic plants, plant cells or plant tissue comprising the introduction of said nucleic acid sequence or said vector into the genome of said plant, plant cell or plant tissue, plant cell transformed with said vector or comprising said DNA sequence, plant obtained by said method;

2. Claims: 1-3, 6-26 partially and 5 completely

At least partially purified protein capable of modulating DNA replication in plants, at least comprising a) one or more of the amino acid sequences of SEQ ID NOS: 6, 7, 10 and 12, b) one or more amino acid sequences having at least 50% amino acid identity with those of a), said protein comprising one or more of the amino acid sequences according to b) having an amino acid identity of at least 90%, said protein having the amino acid sequence as given in SEQ ID NOS: 5, 11 or 13 or having at least 50% amino acid identity

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

with said sequences, said protein being a plant CDC27 protein or a functional analogue thereof, mutein of said protein, peptide comprising a) one or more of the amino acid sequences of SEQ ID NOS: 6, 7, 10 and 12, b) one or more amino acid sequences having at least 50% amino acid identity with those of a), antibody specifically recognizing said protein, said mutein or said peptide, non-genomic DNA sequence coding for said protein, said mutein or said peptide or DNA sequence having a sequence homology of at least 75% of said sequence or complementary DNA sequence thereof, said DNA sequence comprising the DNA sequence as given by SEQ ID NOS: 9, 14 or 15 or having a sequence homology with said sequences of at least 75% or complementary sequence thereof, DNA sequence coding for said peptide corresponding to nucleotides 109-181, 2125-2181 or 1029-1061 of SEQ ID NO: 9, to nucleotides 109-181 or 2092-2148 of SEQ ID NO: 14 or to nucleotides 1-483 of SEQ ID NO: 15 or a DNA sequence having a sequence homology of at least 75% to said sequences or complementary sequence thereof, DNA vector comprising said DNA sequence, method for modulating DNA replication in plant cells, plant parts or plants by conferring the capacity to provide said protein or said mutein in an amount sufficient to modulate DNA replication and/or block mitosis of said cells, method for identifying and/or obtaining proteins capable of modulating the DNA replication in plants comprising a two-hybrid screening assay using CDC27 polynucleotide sequences as bait and a cDNA library of a cell suspension culture as prey, method for the production of transgenic plants, plant cells or plant tissue comprising the introduction of said nucleic acid sequence or said vector into the genome of said plant, plant cell or plant tissue, plant cell transformed with said vector or comprising said DNA sequence, plant obtained by said method;

INTERNATIONAL SEARCH REPORT

International Application No
P 00/06401

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9631608 A	10-10-1996	AU 703644 B	25-03-1999
		AU 2415495 A	05-12-1995
		AU 703525 B	25-03-1999
		AU 5282396 A	23-10-1996
		CA 2188562 A	23-11-1995
		CA 2216406 A	10-10-1996
		CN 1190439 A	12-08-1998
		EP 0759086 A	26-02-1997
		EP 0819174 A	21-01-1998
		WO 9531564 A	23-11-1995
		JP 10500010 T	06-01-1998
		JP 11503319 T	26-03-1999
WO 9841642 A	24-09-1998	AU 6730198 A	12-10-1998
		EP 0972060 A	19-01-2000

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)